

IGF1R (Phospho Tyr1161) Antibody

| | |
|------------------------------|--|
| Catalog No : | YP1219 |
| Reactivity : | Human;Mouse;Rat |
| Applications : | WB;ELISA |
| Target : | IGF1R |
| Fields : | >>EGFR tyrosine kinase inhibitor resistance;>>Endocrine resistance;>>MAPK signaling pathway;>>Ras signaling pathway;>>Rap1 signaling pathway;>>HIF-1 signaling pathway;>>FoxO signaling pathway;>>Oocyte meiosis;>>Autophagy - animal;>>Endocytosis;>>mTOR signaling pathway;>>PI3K-Akt signaling pathway;>>AMPK signaling pathway;>>Longevity regulating pathway;>>Longevity regulating pathway - multiple species;>>Focal adhesion;>>Adherens junction;>>Signaling pathways regulating pluripotency of stem cells;>>Long-term depression;>>Ovarian steroidogenesis;>>Progesterone-mediated oocyte maturation;>>Pathways in cancer;>>Transcriptional misregulation in cancer;>>Proteoglycans in cancer;>>Glioma;>>Prostate cancer;>>Melanoma;>>Breast cancer;>>Hepatocellular carcinoma |
| Gene Name : | IGF1R |
| Protein Name : | Insulin-like growth factor 1 receptor (EC 2.7.10.1) (Insulin-like growth factor I receptor) (IGF-I receptor) (CD antigen CD221) [Cleaved into: Insulin-like growth factor 1 receptor alpha chain; Insuli |
| Human Gene Id : | 3480 |
| Human Swiss Prot No : | P08069 |
| Mouse Gene Id : | 16001 |
| Mouse Swiss Prot No : | Q60751 |
| Rat Gene Id : | 25718 |
| Rat Swiss Prot No : | P24062 |
| Immunogen : | Synthesized phospho derived from human IGF1R (Phospho-Tyr1161) |


| | |
|-------------------------------|--|
| Specificity : | <u>This detects endogenous levels of IGF1R (Phospho-Tyr1161)</u> |
| Formulation : | <u>Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.</u> |
| Source : | <u>Polyclonal, Rabbit,IgG</u> |
| Dilution : | <u>WB 1:500-2000, ELISA 1:10000-20000</u> |
| Purification : | <u>The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.</u> |
| Concentration : | <u>1 mg/ml</u> |
| Storage Stability : | <u>-15°C to -25°C/1 year(Do not lower than -25°C)</u> |
| Observed Band : | <u>pro: 155kD, recetor beta: 95kD</u> |
| Cell Pathway : | <u>Oocyte meiosis;Endocytosis;Focal adhesion;Adherens_Junction;Long-term depression;Progesterone-mediated oocyte maturation;Pathways in cancer;Colorectal cancer;Glioma;Prostate cancer;Melanoma;</u> |
| Background : | <u>This receptor binds insulin-like growth factor with a high affinity. It has tyrosine kinase activity. The insulin-like growth factor I receptor plays a critical role in transformation events. Cleavage of the precursor generates alpha and beta subunits. It is highly overexpressed in most malignant tissues where it functions as an anti-apoptotic agent by enhancing cell survival. Alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. [provided by RefSeq, May 2014],</u> |
| Function : | <u>catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate.,disease:Defects in IGF1R may be a cause in some cases of resistance to insulin-like growth factor 1 (IGF1 resistance) [MIM:270450]. IGF1 resistance is a growth deficiency disorder characterized by intrauterine growth retardation and poor postnatal growth accompanied with increased plasma IGF1.,enzyme regulation:Autophosphorylation activates the kinase activity.,function:This receptor binds insulin-like growth factor 1 (IGF1) with a high affinity and IGF2 with a lower affinity. It has a tyrosine-protein kinase activity, which is necessary for the activation of the IGF1-stimulated downstream signaling cascade. When present in a hybrid receptor with INSR, binds IGF1. PubMed:12138094 shows that hybrid receptors composed of IGF1R and INSR isoform Long are activated with a high affinity by IGF1, with low a</u> |
| Subcellular Location : | <u>Cell membrane ; Single-pass type I membrane protein .</u> |
| Expression : | <u>Found as a hybrid receptor with INSR in muscle, heart, kidney, adipose tissue, skeletal muscle, hepatoma, fibroblasts, spleen and placenta (at protein level).</u> |

Expressed in a variety of tissues. Overexpressed in tumors, including melanomas, cancers of the colon, pancreas prostate and kidney.

| | |
|------------------------|------------|
| Tag : | orthogonal |
| Sort : | 8341 |
| No4 : | 1 |
| Host : | Rabbit |
| Modifications : | Phospho |

Products Images

1 2 3 4



170
130
100
70
55
40
35
25

1 customer's
2 3T3
3 CAC02
4 mouse-liver

Western blot analysis of various lysate, antibody was diluted at 1000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000