

## Gab 1 (phospho Tyr307) Polyclonal Antibody

<b>Catalog No :</b>	YP1070
<b>Reactivity :</b>	Human;Mouse
<b>Applications :</b>	IHC;IF;ELISA
<b>Target :</b>	Gab 1
<b>Fields :</b>	>>EGFR tyrosine kinase inhibitor resistance;>>ErbB signaling pathway;>>Ras signaling pathway;>>Phospholipase D signaling pathway;>>Neurotrophin signaling pathway;>>Bacterial invasion of epithelial cells;>>Proteoglycans in cancer;>>Renal cell carcinoma;>>Hepatocellular carcinoma;>>Gastric cancer
<b>Gene Name :</b>	GAB1
<b>Protein Name :</b>	GRB2-associated-binding protein 1
<b>Human Gene Id :</b>	2549
<b>Human Swiss Prot No :</b>	Q13480
<b>Mouse Gene Id :</b>	14388
<b>Mouse Swiss Prot No :</b>	Q9QYY0
<b>Immunogen :</b>	Synthesized phospho-peptide around the phosphorylation site of human Gab 1 (phospho Tyr307)
<b>Specificity :</b>	Phospho-Gab 1 (Y307) Polyclonal Antibody detects endogenous levels of Gab 1 protein only when phosphorylated at Y307.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	IHC 1:100 - 1:300. ELISA: 1:40000.. IF 1:50-200
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-

---

chromatography using epitope-specific immunogen.

---

**Concentration :** 1 mg/ml

---

**Storage Stability :** -15°C to -25°C/1 year(Do not lower than -25°C)

---

**Molecularweight :** 77kD

---

**Cell Pathway :** ErbB\_HER;Neurotrophin;Renal cell carcinoma;

---

**Background :** GRB2 associated binding protein 1(GAB1) Homo sapiens The protein encoded by this gene is a member of the IRS1-like multisubstrate docking protein family. It is an important mediator of branching tubulogenesis and plays a central role in cellular growth response, transformation and apoptosis. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Aug 2008],

---

**Function :** function:Probably involved in EGF and insulin receptor signaling.,PTM:Phosphorylated on tyrosine residue(s) by the epidermal growth factor receptor (EGFR) and the insulin receptor (INSR). Tyrosine phosphorylation of GAB1 mediates interaction with several proteins that contain SH2 domains.,similarity:Belongs to the GAB family.,similarity:Contains 1 PH domain.,subunit:Interacts with GRB2 and with other SH2-containing proteins. Interacts with phosphorylated LAT2.,

---

**Subcellular Location :** cytosol,

---

**Expression :** Testis,Whole embryo,

---

**Sort :** 6345

---

**No2 :** 3234T

---

**No4 :** 1

---


**Host :** Rabbit

---

**Modifications :** Phospho

---

**Products Images**



Immunohistochemical analysis of paraffin-embedded human Colon cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).