

**c-Myb (phospho Ser532) Polyclonal Antibody**


|                              |   |
|------------------------------|---|
| <b>Catalog No :</b>          | YP1102  |
| <b>Reactivity :</b>          | Human;Mouse   |
| <b>Applications :</b>        | IHC;IF;ELISA  |
| <b>Target :</b>              | c-Myb   |
| <b>Fields :</b>              | >>PI3K-Akt signaling pathway  |
| <b>Gene Name :</b>           | MYB   |
| <b>Protein Name :</b>        | Transcriptional activator Myb   |
| <b>Human Gene Id :</b>       | 4602  |
| <b>Human Swiss Prot No :</b> | P10242  |
| <b>Mouse Gene Id :</b>       | 17863   |
| <b>Mouse Swiss Prot No :</b> | P06876  |
| <b>Immunogen :</b>           | The antiserum was produced against synthesized peptide derived from human Myb around the phosphorylation site of Ser532. AA range:496-545 |
| <b>Specificity :</b>         | Phospho-c-Myb (S532) Polyclonal Antibody detects endogenous levels of c-Myb protein only when phosphorylated at S532.                     |
| <b>Formulation :</b>         | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| <b>Source :</b>              | Polyclonal, Rabbit,IgG  |
| <b>Dilution :</b>            | IHC 1:100 - 1:300. ELISA: 1:5000.. IF 1:50-200  |
| <b>Purification :</b>        | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.                     |
| <b>Concentration :</b>       | 1 mg/ml   |

---


|                               |  |
|-------------------------------|--|
| <b>Storage Stability :</b>    | -15°C to -25°C/1 year(Do not lower than -25°C)   |
| <b>Molecularweight :</b>      | 72kD   |
| <b>Cell Pathway :</b>         | Akt_PKB  |
| <b>Background :</b>           | This gene encodes a protein with three HTH DNA-binding domains that functions as a transcription regulator. This protein plays an essential role in the regulation of hematopoiesis. This gene may be aberrantly expressed or rearranged or undergo translocation in leukemias and lymphomas, and is considered to be an oncogene. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2016],   |
| <b>Function :</b>             | domain:Comprised of 3 domains; an N-terminal DNA-binding domain, a centrally located transcriptional activation domain and a C-terminal domain involved in transcriptional repression.,function:Transcriptional activator; DNA-binding protein that specifically recognize the sequence 5'-YAAC[GT]G-3'. Plays an important role in the control of proliferation and differentiation of hematopoietic progenitor cells.,PTM:Phosphorylated by NLK on multiple sites, which induces proteasomal degradation.,PTM:Ubiquitinated; mediated by SIAH1 and leading to its subsequent proteasomal degradation.,similarity:Contains 3 HTH myb-type DNA-binding domains.,subunit:Binds MYBBP1A. Interacts with HIPK2, MAF and NLK., |
| <b>Subcellular Location :</b> | Nucleus .  |
| <b>Expression :</b>           | Liver,Placenta,Testis,   |
| <b>Sort :</b>                 | 4314   |
| <b>No4 :</b>                  | 1  |
| <b>Host :</b>                 | Rabbit   |
| <b>Modifications :</b>        | Phospho  |

---

## Products Images



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using Myb (Phospho-Ser532) Antibody



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using Myb (Phospho-Ser532) Antibody. The picture on the right is blocked with the phospho peptide.