

BMAL1 (Phospho Ser42) rabbit pAb

Catalog No :	YP1278
Reactivity :	Human;Mouse;Rat
Applications :	WB
Target :	BMAL1
Fields :	>>Circadian rhythm;>>Dopaminergic synapse
Gene Name :	ARNTL BHLHE5 BMAL1 MOP3 PASD3
Protein Name :	BMAL1 (Ser42)
Human Gene Id :	406
Human Swiss Prot No :	O00327
Mouse Gene Id :	11865
Mouse Swiss Prot No :	Q9WTL8
Rat Gene Id :	29657
Rat Swiss Prot No :	Q9EPW1
Immunogen :	Synthesized phospho peptide around human BMAL1 (Ser42)
Specificity :	This antibody detects endogenous levels of Human Mouse Rat BMAL1 (phospho-Ser42)
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:1000-2000

Purification :	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	69kD
Cell Pathway :	Circadian rhythm;
Background :	<p>The protein encoded by this gene is a basic helix-loop-helix protein that forms a heterodimer with CLOCK. This heterodimer binds E-box enhancer elements upstream of Period (PER1, PER2, PER3) and Cryptochrome (CRY1, CRY2) genes and activates transcription of these genes. PER and CRY proteins heterodimerize and repress their own transcription by interacting in a feedback loop with CLOCK/ARNTL complexes. Defects in this gene have been linked to infertility, problems with gluconeogenesis and lipogenesis, and altered sleep patterns. Several transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2014],</p>
Function :	<p>alternative products:Additional isoforms seem to exist,function:ARNTL-CLOCK heterodimers activate E-box element (3'-CACGTG-5') transcription of a number of proteins of the circadian clock. This transcription is inhibited in a feedback loop by PER, and also by CRY proteins.,miscellaneous:CLOCK-ARNTL double mutations within the PAS domains result in synergistic desensitization to high levels of CRY on repression of CLOCK-ARNTL transcriptional activity of PER1 and, disrupt circadian rhythmicity.,PTM:Acetylated on Lys-538 upon dimerization with CLOCK. Acetylation facilitates CRY1-mediated repression.,PTM:Phosphorylated upon dimerization with CLOCK.,PTM:Sumoylated on Lys-259 upon dimerization with CLOCK.,similarity:Contains 1 basic helix-loop-helix (bHLH) domain.,similarity:Contains 1 PAC (PAS-associated C-terminal) domain.,similarity:Contains 2 PAS (PER-ARNT-SIM) domains.,subunit:Component</p>
Subcellular Location :	Nucleus . Cytoplasm . Nucleus, PML body . Shuttles between the nucleus and the cytoplasm and this nucleocytoplasmic shuttling is essential for the nuclear accumulation of CLOCK, target gene transcription and the degradation of the CLOCK-ARNTL/BMAL1 heterodimer. The sumoylated form localizes in the PML body. Sequestered to the cytoplasm in the presence of ID2. .
Expression :	Hair follicles (at protein level). Highly expressed in the adult brain, skeletal muscle and heart.
Sort :	2784
	1

Host: Rabbit**Modifications :** Phospho

Products Images