

eIF4G (Phospho Ser1108) rabbit pAb

Catalog No :	YP1328
Reactivity :	Human;Mouse;Rat
Applications :	WB
Target :	eIF4G
Fields :	>>Viral myocarditis
Gene Name :	EIF4G1 EIF4F EIF4G EIF4GI
Protein Name :	eIF4G (Ser1108)
Human Gene Id :	1981
Human Swiss Prot No :	Q04637
Mouse Gene Id :	208643
Mouse Swiss Prot No :	Q6NZJ6
Immunogen :	Synthesized phospho peptide around human eIF4G (Ser1108)
Specificity :	This antibody detects endogenous levels of Human Mouse Rat eIF4G (phospho-Ser1108)
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:1000-2000
Purification :	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Concentration :	1 mg/ml

Storage Stability : -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 180kD

Cell Pathway : Viral myocarditis;

Background : The protein encoded by this gene is a component of the multi-subunit protein complex EIF4F. This complex facilitates the recruitment of mRNA to the ribosome, which is a rate-limiting step during the initiation phase of protein synthesis. The recognition of the mRNA cap and the ATP-dependent unwinding of 5'-terminal secondary structure is catalyzed by factors in this complex. The subunit encoded by this gene is a large scaffolding protein that contains binding sites for other members of the EIF4F complex. A domain at its N-terminus can also interact with the poly(A)-binding protein, which may mediate the circularization of mRNA during translation. Alternative splicing results in multiple transcript variants, some of which are derived from alternative promoter usage. [provided by RefSeq, Aug 2010],

Function : function:Component of the protein complex eIF4F, which is involved in the recognition of the mRNA cap, ATP-dependent unwinding of 5'-terminal secondary structure and recruitment of mRNA to the ribosome.,PTM:Following infection by certain enteroviruses, rhinoviruses and aphthoviruses, EIF4G1 is cleaved by the viral protease 2A, or the leader protease in the case of aphthoviruses. This shuts down the capped cellular mRNA transcription.,PTM:Phosphorylated at multiple sites in vivo.,sequence caution:Aberrant splicing.,similarity:Belongs to the eIF4G family.,similarity:Contains 1 MI domain.,similarity:Contains 1 MIF4G domain.,similarity:Contains 1 W2 domain.,subunit:eIF4F is a multi-subunit complex, the composition of which varies with external and internal environmental conditions. It is composed of at least EIF4A, EIF4E and EIF4G1/EIF4G3. Interacts with eIF3, mutually exclusive with EIF4A1

Subcellular Location : Cytoplasm, Stress granule .

Expression : Brain,Endometrial tumor,Epithelium,Pancreas,Placent

Sort : 5488

No4 : 1

Host : Rabbit

Modifications : Phospho

Products Images