

PIP5KIII (phospho Ser307) Polyclonal Antibody

Catalog No: YP1162

Reactivity: Human; Mouse; Rat

Applications: IHC;IF;ELISA

Target: PIP5K

Fields: >>Inositol phosphate metabolism;>>Metabolic pathways;>>Phosphatidylinositol

signaling system;>>Phagosome;>>Regulation of actin cytoskeleton

Gene Name: PIKFYVE

Protein Name: 1-phosphatidylinositol 3-phosphate 5-kinase

Q9Y2I7

Q9Z1T6

Human Gene Id: 200576

Human Swiss Prot

No:

Mouse Gene Id: 18711

Mouse Swiss Prot

No:

Immunogen: The antiserum was produced against synthesized peptide derived from human

PIP5K around the phosphorylation site of Ser307. AA range:273-322

Specificity: Phospho-PIP5KIII (S307) Polyclonal Antibody detects endogenous levels of

PIP5KIII protein only when phosphorylated at S307.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution: IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other

applications.

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.



Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight: 237kD

Cell Pathway: Inositol phosphate metabolism; Phosphatidylinositol signaling

system; Endocytosis; Fc gamma R-mediated phagocytosis; Regulates Actin and

Cytoskeleton;

Background: Phosphorylated derivatives of phosphatidylinositol (PtdIns) regulate cytoskeletal

functions, membrane trafficking, and receptor signaling by recruiting protein complexes to cell- and endosomal-membranes. Humans have multiple PtdIns proteins that differ by the degree and position of phosphorylation of the inositol

ring. This gene encodes an enzyme (PIKfyve; also known as

phosphatidylinositol-3-phosphate 5-kinase type III or PIPKIII) that phosphorylates the D-5 position in PtdIns and phosphatidylinositol-3-phosphate (PtdIns3P) to

make PtdIns5P and PtdIns(3,5)biphosphate. The D-5 position also can be phosphorylated by type I PtdIns4P-5-kinases (PIP5Ks) that are encoded by distinct genes and preferentially phosphorylate D-4 phosphorylated PtdIns. In contrast, PIKfyve preferentially phosphorylates D-3 phosphorylated PtdIns. In

addition to being a lipid kinase, PIKf

Function: catalytic activity:ATP + 1-phosphatidyl-1D-myo-inositol 4-phosphate = ADP +

1-phosphatidyl-1D-myo-inositol 4,5-bisphosphate.,disease:Defects in PIKFYVE are the cause of corneal fleck dystrophy (CFD) [MIM:121850]. CFD is an autosomal dominant disorder of the cornea characterized by numerous small white flecks scattered in all levels of the stroma. Although CFD may occasionally cause mild photophobia, patients are typically asymptomatic and have normal vision.,function:Supports the intracellular PIP pool and to a lesser extent, the PI 4,5-P(2) pool. It generates PIP from PI and, to a lesser extent, PI 4,5-P(2) from PI 4-P. There are indications that it phosphorylates the D-5 rather than the D-4 position. Has a role in endosome-related membrane trafficking.,similarity:Contains

1 DEP domain., similarity: Contains 1 FYVE-type zinc finger., similarity: Contains 1

PI5K domain..subcellular location:

Subcellular Location:

Endosome membrane; Peripheral membrane protein. Early endosome membrane; Peripheral membrane protein. Cytoplasmic vesicle, phagosome membrane; Peripheral membrane protein. Late endosome membrane; Peripheral membrane protein. Mainly associated with membranes of the late

endocytic pathway...

Expression: Brain, Epithelium, PCR rescued clones, T-cell,

Sort : 12714


No4: 1




Host: Rabbit

Modifications: Phospho


Products Images



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using PIP5K (Phospho-Ser307) Antibody



Immunofluorescence analysis of HeLa cells, using PIP5K (Phospho-Ser307) Antibody. The picture on the right is blocked with the phospho peptide.



Immunohistochemistry analysis of paraffin-embedded human brain, using PIP5K (Phospho-Ser307) Antibody. The picture on the right is blocked with the phospho peptide.