

**NAGK (Phospho Tyr205) rabbit pAb**

|                              |  |
|------------------------------|--|
| <b>Catalog No :</b>          | YP1626   |
| <b>Reactivity :</b>          | Human;Mouse;Rat  |
| <b>Applications :</b>        | WB;ELISA   |
| <b>Target :</b>              | NAGK   |
| <b>Fields :</b>              | >>Amino sugar and nucleotide sugar metabolism;>>Metabolic pathways;>>Biosynthesis of nucleotide sugars |
| <b>Gene Name :</b>           | NAGK   |
| <b>Protein Name :</b>        | NAGK (phospho-Tyr205)  |
| <b>Human Gene Id :</b>       | 55577  |
| <b>Human Swiss Prot No :</b> | Q9UJ70   |
| <b>Mouse Gene Id :</b>       | 56174  |
| <b>Mouse Swiss Prot No :</b> | Q9QZ08   |
| <b>Rat Gene Id :</b>         | 297393   |
| <b>Rat Swiss Prot No :</b>   | P81799   |
| <b>Immunogen :</b>           | Synthesized peptide derived from human NAGK (phospho-Tyr205)   |
| <b>Specificity :</b>         | This antibody detects endogenous levels of Human,Mouse,Rat NAGK (phospho-Tyr205)                       |
| <b>Formulation :</b>         | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.                                |
| <b>Source :</b>              | Polyclonal, Rabbit,IgG   |
| <b>Dilution :</b>            | WB 1:1000-2000 ELISA 1:5000-20000  |

|                            |   |
|----------------------------|---|
| <b>Purification :</b>      | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. |
| <b>Concentration :</b>     | 1 mg/ml   |
| <b>Storage Stability :</b> | -15°C to -25°C/1 year(Do not lower than -25°C)  |
| <b>Molecularweight :</b>   | 38kD  |
| <b>Expression :</b>        | Ubiquitous.   |
| <b>Sort :</b>              | 10554   |
| <b>No4 :</b>               | 1   |
| <b>Host :</b>              | Rabbit  |
| <b>Modifications :</b>     | Phospho   |

## Products Images