

**eIF2B (Phospho Ser67) rabbit pAb**

<b>Catalog No :</b>	YP1615
<b>Reactivity :</b>	Human;Rat;Mouse;
<b>Applications :</b>	WB;ELISA
<b>Target :</b>	eIF2B
<b>Gene Name :</b>	EIF2S2 EIF2B
<b>Protein Name :</b>	eIF2B (Phospho Ser67)
<b>Human Gene Id :</b>	8894
<b>Human Swiss Prot No :</b>	P20042
<b>Mouse Gene Id :</b>	67204
<b>Mouse Swiss Prot No :</b>	Q99L45
<b>Immunogen :</b>	Synthesized peptide derived from human eIF2B (Phospho Ser67)
<b>Specificity :</b>	This antibody detects endogenous levels of Human eIF2B (Phospho Ser67)
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:1000-2000 ELISA 1:5000-20000
<b>Purification :</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)

**Observed Band :** 69kD

---

**Background :** eukaryotic translation initiation factor 2 subunit beta(EIF2S2) Homo sapiens  
Eukaryotic translation initiation factor 2 (EIF-2) functions in the early steps of protein synthesis by forming a ternary complex with GTP and initiator tRNA and binding to a 40S ribosomal subunit. EIF-2 is composed of three subunits, alpha, beta, and gamma, with the protein encoded by this gene representing the beta subunit. The beta subunit catalyzes the exchange of GDP for GTP, which recycles the EIF-2 complex for another round of initiation. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2015],

---

**Function :** function:eIF-2 functions in the early steps of protein synthesis by forming a ternary complex with GTP and initiator tRNA. This complex binds to a 40S ribosomal subunit, followed by mRNA binding to form a 43S preinitiation complex. Junction of the 60S ribosomal subunit to form the 80S initiation complex is preceded by hydrolysis of the GTP bound to eIF-2 and release of an eIF-2-GDP binary complex. In order for eIF-2 to recycle and catalyze another round of initiation, the GDP bound to eIF-2 must exchange with GTP by way of a reaction catalyzed by eIF-2B.,similarity:Belongs to the eIF-2-beta/eIF-5 family.,subunit:Heterotrimer composed of an alpha, a beta and a gamma chain. Component of an EIF2 complex at least composed of CUGBP1, CALR, CALR3, EIF2S1, EIF2S2, HSP90B1 and HSPA5.,

---

**Sort :** 5462

---

**No4 :** 1

---

**Host :** Rabbit

---

**Modifications :** Phospho

---

## Products Images