

DAPK1 (Phospho Ser289) Rabbit pAb

Catalog No: YP1813

Reactivity: Human; Mouse

Applications: IHC;WB

Target: DAPK1

Fields: >>Autophagy - animal;>>Pathways in cancer;>>Bladder cancer

Gene Name: DAPK1 DAPK

Protein Name: Death-associated protein kinase 1 (DAP kinase 1) (EC 2.7.11.1)

Sequence: P53355

Human Gene Id: 1612

Human Swiss Prot

No:

Mouse Gene ld: 69635

Mouse Swiss Prot

No:

Immunogen: Synthesized peptide derived from human DAPK1 (Phospho Ser289)

Specificity: This antibody detects endogenous levels of DAPK1 (Phospho Ser289) Rabbit

pAb at Human, Mouse

Formulation: Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.

Source: Rabbit,polyclonal

Dilution: WB 1:500-2000 IHC 1:50-200

P53355

Q80YE7

Purification: The antibody was affinity-purified from rabbit serum by affinity-chromatography

using specific immunogen.



Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight: 157kD

Background: death associated protein kinase 1(DAPK1) Homo sapiens Death-associated

protein kinase 1 is a positive mediator of gamma-interferon induced programmed cell death. DAPK1 encodes a structurally unique 160-kD calmodulin dependent serine-threonine kinase that carries 8 ankyrin repeats and 2 putative P-loop consensus sites. It is a tumor suppressor candidate. Alternative splicing results in

multiple transcript variants. [provided by RefSeq, Dec 2013],

Function: catalytic activity:ATP + a protein = ADP + a

phosphoprotein.,cofactor:Magnesium.,enzyme regulation:Negatively regulated by

autophosphorylation on Ser-308.,function:Calcium/calmodulin-dependent

serine/threonine kinase which acts as a positive regulator of

apoptosis.,induction:Up-regulated following treatment with interferon-

gamma.,similarity:Belongs to the protein kinase superfamily.,similarity:Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family. DAP kinase subfamily.,similarity:Contains 1 death domain.,similarity:Contains 1 protein kinase domain.,similarity:Contains 10 ANK repeats.,subcellular location:Colocalizes with

the actin filament system.,

Subcellular [Isoform 1]: Cytoplasm. Cytoplasm, cytoskeleton. Colocalizes with MAP1B in the

Location : microtubules and cortical actin fibers.; [Isoform 2]: Cytoplasm, Cytoplasm,

cytoskeleton.

Expression: Isoform 2 is expressed in normal intestinal tissue as well as in colorectal

carcinomas.

Tag: orthogonal


Sort : 999

No4:

Products Images

2/3





Western Blot analysis of 1 HeLa cell, 2 LPS 100ng/mL 30min treated ,using primary antibody at 1:1000 dilution. Secondary antibody(catalog#:RS23920) was diluted at 1:10000