

## USF1 (Phospho Thr153) rabbit pAb

<b>Catalog No :</b>	YP1746
<b>Reactivity :</b>	Human;Mouse;Rat
<b>Applications :</b>	WB
<b>Target :</b>	USF1
<b>Gene Name :</b>	USF1 BHLHB11 USF
<b>Protein Name :</b>	USF1 (Phospho-Thr153)
<b>Human Gene Id :</b>	7391
<b>Human Swiss Prot No :</b>	P22415
<b>Mouse Gene Id :</b>	22278
<b>Mouse Swiss Prot No :</b>	Q61069
<b>Immunogen :</b>	Synthesized peptide derived from human USF1 (Phospho-Thr153)
<b>Specificity :</b>	This antibody detects endogenous levels of USF1 (Phospho-Thr153) at Human, Mouse,Rat
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500-2000
<b>Purification :</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)

---

**Molecularweight :** 34kD

---

**Background :** This gene encodes a member of the basic helix-loop-helix leucine zipper family, and can function as a cellular transcription factor. The encoded protein can activate transcription through pyrimidine-rich initiator (Inr) elements and E-box motifs. This gene has been linked to familial combined hyperlipidemia (FCHL). Alternative splicing of this gene results in multiple transcript variants. A related pseudogene has been defined on chromosome 21. [provided by RefSeq, Feb 2013],

---

**Function :** disease:Genetic variations in USF1 are associated with combined hyperlipidemia type 1 (HYPLIP1) [MIM:602491]; also known as familial combined hyperlipidemia type 1 (FCHL1). HYPLIP1 is characterized by elevated levels of serum total cholesterol, triglycerides or both, and is observed in about 20% of individuals with premature coronary heart disease.,function:Transcription factor that binds to a symmetrical DNA sequence (E-boxes) (5'-CACGTG-3') that is found in a variety of viral and cellular promoters.,similarity:Contains 1 basic helix-loop-helix (bHLH) domain.,subunit:Efficient DNA binding requires dimerization with another bHLH protein. Binds DNA as an homodimer or a heterodimer (USF1/USF2). Interacts with varicella-zoster virus IE62 protein.,

---

**Subcellular Location :** Nucleus.

---

**Expression :** Kidney,

---

**Tag :** orthogonal

---

**Sort :** 25225

---

**No4 :** 1

---


**Host :** Rabbit

---

**Modifications :** Phospho

---

**Products Images**



Western Blot analysis of 1 HeLa cell, 2 LPS 100ng/mL 30min treated ,using primary antibody at 1:1000 dilution. Secondary antibody(catalog#:RS23920) was diluted at 1:10000