

GATA-2/3 (Acetyl Lys336/304) Polyclonal Antibody

Catalog No :	YK0076
Reactivity :	Human:K336/304;Mouse:K336/304;Rat:K336/305
Applications :	WB;ELISA
Target :	GATA-2/3
Gene Name :	GATA2/GATA3
Protein Name :	GATA binding protein 2/GATA binding protein 3
Human Gene Id :	2624
Human Swiss Prot No :	P23769/P23771
Immunogen :	Synthesized acetyl-peptide from human protein at AA range: 320-390
Specificity :	This antibody detects endogenous levels of GATA-2/3 at Human:K336/304;Mouse:K336/304;Rat:K336/305, It doesn't react with total protein.
Formulation :	PBS, pH 7.4, containing 0.02% sodium azide as Preservative and 50% Glycerol.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-10000 ELISA: 1:10000
Purification :	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	50kD
Background :	This gene encodes a member of the GATA family of zinc-finger transcription

factors that are named for the consensus nucleotide sequence they bind in the promoter regions of target genes. The encoded protein plays an essential role in regulating transcription of genes involved in the development and proliferation of hematopoietic and endocrine cell lineages. Alternative splicing results in multiple transcript variants.[provided by RefSeq, Mar 2009],

Function :

function:Transcriptional activator which regulates endothelin-1 gene expression in endothelial cells. Binds to the consensus sequence 5'-AGATAG-3',,sequence caution:Several sequencing errors.,similarity:Contains 2 GATA-type zinc fingers.,tissue specificity:Endothelial cells.,

Subcellular Location :

Nucleus.

Expression :

Endothelial cells.

Sort :

6479

No4 :

1


Host :

Rabbit

Modifications :

Acetyl

Products Images



Western Blot analysis of hepg2 cells using Antibody diluted at 500. Secondary antibody(catalog#:RS0002) was diluted at 1:20000