

## Histone H3 (Tri-Methyl-K10) Polyclonal Antibody


<b>Catalog No :</b>	YH0012
<b>Reactivity :</b>	Human;Rat;Mouse
<b>Applications :</b>	WB;ELISA
<b>Target :</b>	Histone H3
<b>Fields :</b>	>>Neutrophil extracellular trap formation;>>Alcoholism;>>Shigellosis;>>Transcriptional misregulation in cancer;>>Systemic lupus erythematosus
<b>Gene Name :</b>	Histone H3
<b>Protein Name :</b>	Histone H3
<b>Human Gene Id :</b>	8350
<b>Human Swiss Prot No :</b>	P68431/Q71DI3/P84243/Q6NXT2
<b>Immunogen :</b>	Synthetic Tri-Methyl peptide from human protein at AA range: 10
<b>Specificity :</b>	The antibody detects endogenous Tri-Methyl-Histone H3 (K10)
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500-2000, ELISA 1:10000-20000
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Observed Band :</b>	15kD

<b>Cell Pathway :</b>	Systemic lupus erythematosus;
<b>Background :</b>	<p>Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3. [provided by RefSeq, Aug 2015],</p>
<b>Function :</b>	<p>caution:Was originally (PubMed:2587222) thought to originate from mouse.,developmental stage:Expressed during S phase, then expression strongly decreases as cell division slows down during the process of differentiation.,function:Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.,mass spectrometry:Monoisotopic with N-acetylserine PubMed:16457589,miscellaneous:This histone is only present in mammals and is enriched in acetylation of Lys-15 and dimethylation of Lys-10 (H3K9me2).,PTM:Acetylation is generally I</p>
<b>Subcellular Location :</b>	Nucleus. Chromosome.
<b>Expression :</b>	Blood,Epithelium,Kidney,Lung,Ovary,Spleen,Uterus,
<b>Sort :</b>	7589
<b>No4 :</b>	1
<b>Host :</b>	Rabbit
<b>Modifications :</b>	Methyl

## Products Images

170—  
130—  
100—  
70—  
55—  
40—  
35—  
25—  
15—

1, 3T3 2, mouse-kidney 3, KB  
4, K562 5, HeLa 6, 293T



Western blot analysis of 293T lysate, antibody was diluted at 2000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000