

## FoxO1A (Acetyl Lys248) rabbit pAb

Catalog No: YK0111

**Reactivity:** Human; Mouse; Rat

**Applications:** WB;ELISA

Target: FoxO1

**Fields:** >>FoxO signaling pathway;>>AMPK signaling pathway;>>Longevity regulating

pathway;>>Longevity regulating pathway - multiple species;>>Cellular senescence;>>Insulin signaling pathway;>>Thyroid hormone signaling pathway;>>Glucagon signaling pathway;>>Insulin resistance;>>AGE-RAGE

signaling pathway in diabetic complications;>>Alcoholic liver

disease;>>Shigellosis;>>Human papillomavirus infection;>>Pathways in cancer;>>Transcriptional misregulation in cancer;>>Prostate cancer

Gene Name: FOXO1 FKHR FOXO1A

Protein Name: FoxO1A (Acetyl Lys248)

Q12778

**Q9R1E0** 

Human Gene Id: 2308

**Human Swiss Prot** 

No:

Mouse Gene ld: 56458

**Mouse Swiss Prot** 

No:

Rat Gene ld: 84482

Rat Swiss Prot No: G3V7R4

**Immunogen:** Synthesized peptide derived from human FoxO1A (Acetyl Lys248)

**Specificity:** This antibody detects endogenous levels of Human, Mouse, Rat FoxO1A (Acetyl

Lys248)

**Formulation:** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

1/3



Source: Polyclonal, Rabbit, IgG

**Dilution:** WB 1:1000-2000 ELISA 1:5000-20000

**Purification:** The antibody was affinity-purified from rabbit serum by affinity-chromatography

using specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 72kD

**Background:** disease:Chromosomal aberrations involving FOXO1 are a cause of

rhabdomyosarcoma 2 (RMS2) [MIM:268220]; also known as alveolar rhabdomyosarcoma. Translocation (2;13)(q35;q14) with PAX3; translocation

t(1;13)(p36;q14) with PAX7. The resulting protein is a transcriptional

activator., function: Transcription factor., PTM: Phosphorylated by AKT1; insulin-induced (By similarity). IGF1 rapidly induces phosphorylation of Ser-256, Thr-24, and Ser-319. Phosphorylation of Ser-256 decreases DNA-binding activity and promotes the phosphorylation of Thr-24, and Ser-319, permitting phosphorylation of Ser-322 and Ser-325, probably by CK1, leading to nuclear exclusion and loss of function. Phosphorylation of Ser-329 is independent of IGF1 and leads to

reduced function. Phosphorylated upon DNA damage, probably by ATM or ATR., similarity: Contains 1 fork-head DNA-binding domain., subcellular location: Shuttles between cytoplasm and nucleus., subunit: Interacts with

LRPPRC., tissue specificity: Ubiquitous.,

Function: blood vessel development, vasculature development, transcription, regulation of

transcription, DNA-dependent, regulation of transcription from RNA polymerase II promoter, anti-apoptosis, cell surface receptor linked signal transduction, enzyme linked receptor protein signaling pathway, transmembrane receptor protein tyrosine kinase signaling pathway, insulin receptor signaling pathway, response to endogenous stimulus, response to hormone stimulus, positive regulation of biosynthetic process, response to organic substance, positive regulation of macromolecule biosynthetic process, positive regulation of macromolecule metabolic process, positive regulation of gene expression, regulation of cell death, positive regulation of cellular biosynthetic process, response to insulin stimulus, cellular response to hormone

stimulus, regulation of cell proliferati

Subcellular Location:

Cytoplasm . Nucleus . Shuttles between the cytoplasm and nucleus. Largely nuclear in unstimulated cells (PubMed:11311120, PubMed:12228231,

PubMed:19221179, PubMed:21245099, PubMed:20543840,

PubMed:25009184). In osteoblasts, colocalizes with ATF4 and RUNX2 in the nucleus (By similarity). Serum deprivation increases localization to the nucleus, leading to activate expression of SOX9 and subsequent chondrogenesis (By



similarity). Insulin-induced phosphorylation at Ser-256 by PKB/AKT1 leads, via stimulation of Thr-24 phosphorylation, to binding of 14-3-3 proteins and nuclear export to the cytoplasm where it is degraded by the ubiquitin-proteosomal pathway (PubMed:11237865, PubMed:12228231). Phosphorylation at Ser-249 by CDK1 disrupts binding of 14-3-3 proteins and promotes nuclear accumulation

Expression :	Ubiquitous.
Sort :	6250
No4 :	_1
Host :	Rabbit
Modifications :	Acetyl

## **Products Images**

3/3