

CD38 (PN0604) Nb-FC recombinant antibody

| | |
|------------------------------|--|
| Catalog No : | YA0343 |
| Reactivity : | Human |
| Applications : | ELISA |
| Target : | CD38 |
| Gene Name : | CD38 |
| Protein Name : | ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1 (EC 3.2.2.6) (2'-phospho-ADP-ribosyl cyclase) (2'-phospho-ADP-ribosyl cyclase/2'-phospho-cyclic-ADP-ribose transferase) (EC 2.4.99.20) (2'-phospho-cyc |
| Human Gene Id : | 952 |
| Human Swiss Prot No : | P28907 |
| Immunogen : | Purified recombinant Human CD38 |
| Specificity : | This recombinant monoclonal antibody can detects endogenous levels of CD38 protein. |
| Formulation : | Phosphate-buffered solution |
| Source : | Camel, chimeric fusion of Nanobody (VHH) and mouse IgG1 Fc domain , recombinantly produced from 293F cell |
| Dilution : | ELISA 1:5000-100000 |
| Purification : | Recombinant Expression and Affinity purified |
| Concentration : | Please check the information on the tube |
| Storage Stability : | -15°C to -25°C/1 year(Avoid freeze / thaw cycles) |
| Background : | The protein encoded byThis gene is a non-lineage-restricted, type II transmembrane glycoprotein that synthesizes and hydrolyzes cyclic adenosine |

5'-diphosphate-ribose, an intracellular calcium ion mobilizing messenger. The release of soluble protein and the ability of membrane-bound protein to become internalized indicate both extracellular and intracellular functions for the protein. This protein has an N-terminal cytoplasmic tail, a single membrane-spanning domain, and a C-terminal extracellular region with four N-glycosylation sites. Crystal structure analysis demonstrates that the functional molecule is a dimer, with the central portion containing the catalytic site. It is used as a prognostic marker for patients with chronic lymphocytic leukemia. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2015]

Function :

catalytic activity:NAD(+) + H₂O = ADP-ribose + nicotinamide.,developmental stage:Preferentially expressed at both early and late stages of the B and T-cell maturation. It is also detected on erythroid and myeloid progenitors in bone marrow, where the level of surface expression was shown to decrease during differentiation of blast-forming unit E to colony-forming unit E.,enzyme regulation:ATP inhibits the hydrolyzing activity.,Synthesizes cyclic ADP-ribose, a second messenger for glucose-induced insulin secretion. Also has cADPr hydrolase activity. Also moonlights as a receptor in cells of the immune system.,online information:CD38 entry,similarity:Belongs to the ADP-ribosyl cyclase family.,tissue specificity:Expressed at high levels in pancreas, liver, kidney, brain, testis, ovary, placenta, malignant lymphoma and neuroblastoma.,

Subcellular Location :

Membrane; Single-pass type II membrane protein.

Expression :

Expressed at high levels in pancreas, liver, kidney, brain, testis, ovary, placenta, malignant lymphoma and neuroblastoma.

Tag :

recombinant

Sort :

3573

No4 :

1

Speciality :

Nanobody

Products Images