

## CD223/LAG-3 (PN0567) Nb-FC recombinant antibody

Catalog No: YA0193

Reactivity: Human

**Applications:** ELISA

Target: CD223/LAG-3

Gene Name: LAG3 FDC

Protein Name: Lymphocyte activation gene 3 protein (LAG-3) (CD antigen CD223) [Cleaved

into: Secreted lymphocyte activation gene 3 protein (sLAG-3)]

Human Gene Id: 3902

**Human Swiss Prot** 

No:

P18627

Immunogen:

Purified recombinant Human CD223

**Specificity:** This recombinant monoclonal antibody can detects endogenous levels of

CD223/LAG-3 protein.

Formulation: Phosphate-buffered solution

**Source:** Camel, chimeric fusion of Nanobody (VHH) and mouse IgG1 Fc domain,

recombinantly produced from 293F cell

**Dilution:** ELISA 1:5000-100000

**Purification:** Recombinant Expression and Affinity purified

**Concentration:** Please check the information on the tube

Storage Stability: -15°C to -25°C/1 year(Avoid freeze / thaw cycles)

**Background:** Lymphocyte-activation protein 3 belongs to Ig superfamily and contains 4

extracellular Ig-like domains. The LAG3 gene contains 8 exons. The sequence data, exon/intron organization, and chromosomal localization all indicate a close



relationship of LAG3 to CD4. [provided by RefSeq, Jul 2008]

**Function:** Involved in lymphocyte activation. Binds to HLA class-II

antigens.,similarity:Contains 1 Ig-like V-type (immunoglobulin-like) domain.,similarity:Contains 3 Ig-like C2-type (immunoglobulin-like)

domains., tissue specificity: On cell surface of activated NK and T-lymphocytes.,

Subcellular Location:

[Lymphocyte activation gene 3 protein]: Cell membrane; Single-pass type I membrane protein.; [Secreted lymphocyte activation gene 3 protein]: Secreted.

Produced following cleavage of the main chain. .

**Expression :** Primarily expressed in activated T-cells and a subset of natural killer (NK) cells.

Tag: recombinant

**Sort :** 3471

No3: CD223 nanobody

**No4**:

**Speciality:** Nanobody

## **Products Images**

2/2