

CD25 (PN0488) Nb-FC recombinant antibody

Catalog No :	YA0205
Reactivity :	Human
Applications :	ELISA
Target :	CD25
Gene Name :	IL2RA
Protein Name :	Interleukin-2 receptor subunit alpha (IL-2 receptor subunit alpha) (IL-2-RA) (IL-2R subunit alpha) (IL2-RA) (TAC antigen) (p55) (CD antigen CD25)
Human Gene Id :	3559
Human Swiss Prot No :	P01589
Immunogen :	Purified recombinant Human CD25
Specificity :	This recombinant monoclonal antibody can detects endogenous levels of CD25 protein.
Formulation :	Phosphate-buffered solution
Source :	Camel, chimeric fusion of Nanobody (VHH) and mouse IgG1 Fc domain , recombinantly produced from 293F cell
Dilution :	ELISA 1:5000-100000
Purification :	Recombinant Expression and Affinity purified
Concentration :	Please check the information on the tube
Storage Stability :	-15°C to -25°C/1 year(Avoid freeze / thaw cycles)
Cell Pathway :	Cytokine-cytokine receptor interaction;Endocytosis;Jak_STAT;Hematopoietic cell lineage;

Background :	The interleukin 2 (IL2) receptor alpha (IL2RA) and beta (IL2RB) chains, together with the common gamma chain (IL2RG), constitute the high-affinity IL2 receptor. Homodimeric alpha chains (IL2RA) result in low-affinity receptor, while homodimeric beta (IL2RB) chains produce a medium-affinity receptor. Normally an integral-membrane protein, soluble IL2RA has been isolated and determined to result from extracellular proteolysis. Alternately-spliced IL2RA mRNAs have been isolated, but the significance of each is presently unknown. Mutations in This gene are associated with interleukin 2 receptor alpha deficiency.[provided by RefSeq, Nov 2009]
Function :	disease:Genetic variations in IL2RA are associated with susceptibility to insulin-dependent diabetes mellitus type 10 (IDDM10) [MIM:601942].,Receptor for interleukin-2.,online information:IL2RA mutation db,similarity:Contains 2 Sushi (CCP/SCR) domains.,subunit:Non-covalent dimer of an alpha and a beta chains. IL2R exists in 3 different forms: a high affinity dimer, an intermediate affinity monomer (beta chain), and a low affinity monomer (alpha chain). The high and intermediate affinity forms also associate with a gamma chain.,
Subcellular Location :	Membrane; Single-pass type I membrane protein.
Expression :	Thymus
Tag :	recombinant
Sort :	3498
No4 :	1
Speciality :	Nanobody

Products Images