

**CD269/BCMA (PN0554) Nb-FC recombinant antibody**

<b>Catalog No :</b>	YA0223
<b>Reactivity :</b>	Human
<b>Applications :</b>	FCM;ELISA
<b>Target :</b>	CD269/BCMA
<b>Gene Name :</b>	TNFRSF17 BCM BCMA
<b>Protein Name :</b>	Tumor necrosis factor receptor superfamily member 17 (B-cell maturation protein) (CD antigen CD269)
<b>Human Gene Id :</b>	608
<b>Human Swiss Prot No :</b>	Q02223
<b>Immunogen :</b>	Purified recombinant Human CD269
<b>Specificity :</b>	This recombinant monoclonal antibody can detects endogenous levels of CD269/BCMA protein.
<b>Formulation :</b>	Phosphate-buffered solution
<b>Source :</b>	Camel, chimeric fusion of Nanobody (VHH) and mouse IgG1 Fc domain , recombinantly produced from 293F cell
<b>Dilution :</b>	ELISA 1:5000-100000;FCM 1-2µg/Test
<b>Purification :</b>	Recombinant Expression and Affinity purified
<b>Concentration :</b>	Please check the information on the tube
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Avoid freeze / thaw cycles)
<b>Cell Pathway :</b>	Cytokine-cytokine receptor interaction;Intestinal immune network for IgA production;

**Background :** The protein encoded by This gene is a member of the TNF-receptor superfamily. This receptor is preferentially expressed in mature B lymphocytes, and may be important for B cell development and autoimmune response. This receptor has been shown to specifically bind to the tumor necrosis factor (ligand) superfamily, member 13b (TNFSF13B/TALL-1/BAFF), and to lead to NF-kappaB and MAPK8/JNK activation. This receptor also binds to various TRAF family members, and thus may transduce signals for cell survival and proliferation. [provided by RefSeq, Jul 2008]

---

**Function :** disease: A chromosomal aberration involving TNFRSF17 is found in a form of T-cell acute lymphoblastic leukemia (T-ALL). Translocation t(4;16)(q26;p13) with IL2., Receptor for TNFSF13B/BLyS/BAFF and TNFSF13/APRIL. Promotes B-cell survival and plays a role in the regulation of humoral immunity. Activates NF-kappa-B and JNK., similarity: Contains 1 TNFR-Cys repeat., subcellular location: Perinuclear Golgi-like structures., subunit: Associates with TRAF1, TRAF2, TRAF3, TRAF5 and TRAF6., tissue specificity: Expressed in mature B-cells, but not in T-cells or monocytes.,

---

**Subcellular Location :** Cell membrane; Single-pass type III membrane protein. Endomembrane system; Single-pass type III membrane protein. Perinuclear Golgi-like structures.

---

**Expression :** Expressed in mature B-cells, but not in T-cells or monocytes.

---

**Tag :** recombinant

---

**Sort :** 3502

---

**No4 :** 1

---

**Speciality :** Nanobody

---

## Products Images