

CD70 (PN0448) Nb-FC recombinant antibody

Catalog No: YA0477

Reactivity: Human

Applications: ELISA

Target: CD70

Gene Name: CD70 CD27L CD27LG TNFSF7

Protein Name: CD70 antigen (CD27 ligand) (CD27-L) (Tumor necrosis factor ligand

superfamily member 7) (CD antigen CD70)

Human Gene Id: 970

Human Swiss Prot

No:

Immunogen: Purified recombinant Human CD70

P32970

Specificity: This recombinant monoclonal antibody can detects endogenous levels of CD70

protein.

Formulation: Phosphate-buffered solution

Source: Camel, chimeric fusion of Nanobody (VHH) and mouse IgG1 Fc domain,

recombinantly produced from 293F cell

Dilution: ELISA 1:5000-100000

Purification: Recombinant Expression and Affinity purified

Concentration: Please check the information on the tube

Storage Stability: -15°C to -25°C/1 year(Avoid freeze / thaw cycles)

Cell Pathway: Cytokine-cytokine receptor interaction;

Background: The protein encoded by This gene is a cytokine that belongs to the tumor

1/2

necrosis factor (TNF) ligand family. This cytokine is a ligand for TNFRSF27/CD27. It is a surface antigen on activated, but not on resting, T and B lymphocytes. It induces proliferation of costimulated T cells, enhances the generation of cytolytic T cells, and contributes to T cell activation. This cytokine is also reported to play a role in regulating B-cell activation, cytotoxic function of natural killer cells, and immunoglobulin sythesis. [provided by RefSeq, Jul 2008]

Function: Cytokine that binds to CD27. Plays a role in T-cell activation. Induces the

proliferation of costimulated T-cells and enhances the generation of cytolytic T-

cells., similarity: Belongs to the tumor necrosis factor family., subunit: Homotrimer.,

Subcellular Location:

Membrane; Single-pass type II membrane protein.

Expression: B-cell, Kidney

Tag: recombinant

Sort: 3652

No1: 14-0701-85

No4:

Speciality: Nanobody

Products Images

