

IL-1B (PN0164) Nb-FC recombinant antibody

Catalog No :	YA0609
Reactivity :	Human
Applications :	ELISA
Target :	IL-1B
Gene Name :	IL1B IL1F2
Protein Name :	Interleukin-1 beta (IL-1 beta) (Catabolin)
Human Gene Id :	3553
Human Swiss Prot No :	P01584
Immunogen :	Purified recombinant Human IL-1B
Specificity :	This recombinant monoclonal antibody can detects endogenous levels of IL-1B protein.
Formulation :	Phosphate-buffered solution
Source :	Camel, chimeric fusion of Nanobody (VHH) and mouse IgG1 Fc domain , recombinantly produced from 293F cell
Dilution :	ELISA 1:5000-100000
Purification :	Recombinant Expression and Affinity purified
Concentration :	Please check the information on the tube
Storage Stability :	-15°C to -25°C/1 year(Avoid freeze / thaw cycles)
Background :	IL-1 β in humans and mice does not encode a typical signal peptide and, as a result, newly synthesized pro-IL-1 β accumulates within the cytoplasm of activated monocytes and macrophages . Conversion of the inactive pro-IL-1 β to its mature form requires the proteolytic action of IL-1 β -converting enzyme (ICE), also termed

caspace-1 . Secretion of mature IL-1 β from LPS-activated monocytes/macrophages is not a constitutive process. These cells must encounter a secondary stimulus that specifically activates the posttranslational processing events . Moreover, owing to its pro-inflammatory nature, IL-1 β is regarded as a tumor-promoting cytokine. In fact, enhanced tumor metastasis and angiogenesis has been observed under the influence of IL-1 β . IL-1 β is able to facilitate tumor progression in murine models of lung cancer. In addition, upregulation of metastasis and tumor angiogenesis by IL-1 β has been associated with increased activity of matrix metalloproteinases and expression of the pro-angiogenic molecule hepatocyte growth factor .

Function :

Interleukin (IL)-1 β is a cytokine with a key role in the pathophysiology of local and systemic inflammation. IL-1 β induces cytokine, chemokine, proinflammatory molecule secretion, and adhesion molecule expression in diverse cells.

Subcellular Location :

Cytoplasm, cytosol . Secreted . Lysosome . Secreted, extracellular exosome . The precursor is cytosolic (PubMed:15192144). In response to inflammasome-activating signals, such as ATP for NLRP3 inflammasome or bacterial flagellin for NLRC4 inflammasome, cleaved and secreted (PubMed:24201029, PubMed:33377178, PubMed:33883744). Mature form is secreted and released in the extracellular milieu by passing through the gasdermin-D (GSDMD) pore (PubMed:33883744). In contrast, the precursor form is not released, due to the presence of an acidic region that is proteolytically removed by CASP1 during maturation (PubMed:33883744). The secretion is dependent on protein unfolding and facilitated by the cargo receptor TMED10 (PubMed:32272059). .

Expression :

Expressed in activated monocytes/macrophages (at protein level).

Tag :

recombinant

Sort :

16

No4 :

1

Speciality :

Nanobody

Products Images