

EphB4 Monoclonal Antibody

Catalog No :	YM0232
Reactivity :	Human
Applications :	WB;IHC;IF;ELISA
Target :	EphB4
Fields :	>>Axon guidance
Gene Name :	EPHB4
Protein Name :	Ephrin type-B receptor 4
Human Gene Id :	2050
Human Swiss Prot No :	P54760
Mouse Swiss Prot No :	P54761
Immunogen :	Purified recombinant fragment of EphB4 expressed in E. Coli.
Specificity :	EphB4 Monoclonal Antibody detects endogenous levels of EphB4 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Monoclonal, Mouse
Dilution :	WB 1:500 - 1:2000. IHC 1:200 - 1:1000. ELISA: 1:10000.. IF 1:50-200
Purification :	Affinity purification
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Molecularweight :	108kD
Cell Pathway :	Axon guidance;

P References : 1. J. Chrencik, A. Brooun, M. Recht. Structure. 2006 Feb;14(2):321-30.
2. Qinghua WU, Zhenhe SUO, Bjorn RISBERG. Pathol Oncol Res. 2004;10(1):26-33.

Background : Ephrin receptors and their ligands, the ephrins, mediate numerous developmental processes, particularly in the nervous system. Based on their structures and sequence relationships, ephrins are divided into the ephrin-A (EFNA) class, which are anchored to the membrane by a glycosylphosphatidylinositol linkage, and the ephrin-B (EFNB) class, which are transmembrane proteins. The Eph family of receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. Ephrin receptors make up the largest subgroup of the receptor tyrosine kinase (RTK) family. The protein encoded by this gene binds to ephrin-B2 and plays an essential role in vascular development. [provided by RefSeq, Jul 2008],

Function : catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate.,function:Receptor for members of the ephrin-B family. Binds to ephrin-B2. May have a role in events mediating differentiation and development.,PTM:Autophosphorylated.,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family.,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family. Ephrin receptor subfamily.,similarity:Contains 1 protein kinase domain.,similarity:Contains 1 SAM (sterile alpha motif) domain.,similarity:Contains 2 fibronectin type-III domains.,tissue specificity:Abundantly expressed in placenta and in a range of primary tissues and malignant cell lines. Expressed in fetal, but not adult, brain, and in primitive and myeloid, but not lymphoid, hematopoietic cells.,

Subcellular Location : Cell membrane ; Single-pass type I membrane protein .

Expression : Abundantly expressed in placenta but also detected in kidney, liver, lung, pancreas, skeletal muscle and heart. Expressed in primitive and myeloid, but not lymphoid, hematopoietic cells. Also observed in cell lines derived from liver, breast, colon, lung, melanocyte and cervix.


Sort : 5653

No4 : 1


Host : Mouse

Modifications : Unmodified

Products Images



Western Blot analysis using EphB4 Monoclonal Antibody against truncated EphB4 recombinant protein.



Immunohistochemistry analysis of paraffin-embedded Human pancreas carcinoma (left) and breast carcinoma (right) tissue, showing membrane and cytoplasmic (pancreas carcinoma) localization, membrane (breast carcinoma) localization with DAB staining using Ep