

Calnexin Monoclonal Antibody

Catalog No: YM0089

Reactivity: Human

Applications: WB;IF;ELISA

Target: Calnexin

Fields: >>Protein processing in endoplasmic reticulum;>>Phagosome;>>Antigen

processing and presentation;>>Thyroid hormone synthesis;>>Human T-cell

leukemia virus 1 infection

Gene Name: CANX

Protein Name: Calnexin

Human Gene Id: 821

Human Swiss Prot

No:

Mouse Swiss Prot

No:

Immunogen: Synthetic peptide corresponding to aa (CEAAEERPWLWVVYILTVAL) of human

Calnexin, conjugated to KLH.

Specificity: Calnexin Monoclonal Antibody detects endogenous levels of Calnexin protein.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Monoclonal, Mouse

Dilution: WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other

applications.

P27824

P35564

Purification: Affinity purification

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

1/3



Molecularweight: 68kD

Cell Pathway : Antigen processing and presentation;

P References : 1. Science. 2003 Feb 28;299(5611):1394-7.

2. Exp Cell Res. 2004 Mar 10;294(1):244-53.

3. Science. 2004 Apr 23;304(5670):600-2.

Background: This gene encodes a member of the calnexin family of molecular chaperones.

The encoded protein is a calcium-binding, endoplasmic reticulum

(ER)-associated protein that interacts transiently with newly synthesized N-linked glycoproteins, facilitating protein folding and assembly. It may also play a central role in the quality control of protein folding by retaining incorrectly folded protein subunits within the ER for degradation. Alternatively spliced transcript variants encoding the same protein have been described. [provided by RefSeq, Jul 2008],

Function: function:Calcium-binding protein that interacts with newly synthesized

glycoproteins in the endoplasmic reticulum. It may act in assisting protein assembly and/or in the retention within the ER of unassembled protein subunits. It

seems to play a major role in the quality control apparatus of the ER by the

retention of incorrectly folded proteins.,online information:Calnexin

entry, similarity: Belongs to the calreticulin family., subcellular location: Identified by

mass spectrometry in melanosome fractions from stage I to stage IV.,

Subcellular Location:

Endoplasmic reticulum membrane ; Single-pass type I membrane protein . Endoplasmic reticulum . Melanosome . Identified by mass spectrometry in

melanosome fractions from stage I to stage IV (PubMed:12643545, PubMed:17081065). The palmitoylated form preferentially localizes to the

perinuclear rough ER (PubMed:22314232). .

Expression : B-cell lymphoma, Epithelium, Fibroblast, Keratinocyte, Kidney, Liver, Lymph, Placen

ta,Platelet,


Sort : 3063

No4: 1


Host: Mouse

Modifications: Unmodified

Products Images



Western Blot analysis using Calnexin Monoclonal Antibody against A431 (1), HeLa (2), MCF-7 (3) and A549 (4) cell lysate.



Confocal immunofluorescence analysis of Hela cells using Calnexin Monoclonal Antibody (green).