

CD73 Monoclonal Antibody

Catalog No: YM0133

Reactivity: Human

Applications: IHC;IF;ELISA

Target: CD73

Fields: >>Purine metabolism;>>Pyrimidine metabolism;>>Nicotinate and nicotinamide

metabolism;>>Metabolic pathways;>>Nucleotide metabolism

Gene Name: NT5E

Protein Name: 5'-nucleotidase

Human Gene Id: 4907

Human Swiss Prot P21589

No:

Mouse Swiss Prot

No:

Immunogen: Purified recombinant fragment of CD73 expressed in E. Coli.

Specificity: CD73 Monoclonal Antibody detects endogenous levels of CD73 protein.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Monoclonal, Mouse

Dilution : IHC 1:200 - 1:1000. ELISA: 1:10000.. IF 1:50-200

Purification: Affinity purification

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Cell Pathway: Purine metabolism;Pyrimidine metabolism;Nicotinate and nicotinamide

metabolism;

Q61503

P References:

1. Oncol Rep. 2007 Jun;17(6):1341-6.

2. Neurochem Int. 2003 Dec;43(7):621-8.

Background:

The protein encoded by this gene is a plasma membrane protein that catalyzes the conversion of extracellular nucleotides to membrane-permeable nucleosides. The encoded protein is used as a determinant of lymphocyte differentiation. Defects in this gene can lead to the calcification of joints and arteries. Two transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Mar 2011],

Function:

catalytic activity: A 5'-ribonucleotide + H(2)O = a ribonucleoside + phosphate., cofactor: Zinc., disease: There is a decrease in the activity of NT5 in B-cell chronic lymphocytic leukemia., function: Hydrolyzes extracellular nucleotides into membrane permeable nucleosides., similarity: Belongs to the 5'-nucleotidase family., subunit: Homodimer; disulfide-linked.,

Subcellular Location:

Cell membrane; Lipid-anchor, GPI-anchor.

Expression : Leukocyte, Liver, Placenta, Skin,

Tag: hot


Sort: 3659

No4:

Host: Mouse

Modifications: Unmodified

Products Images



Immunohistochemistry analysis of paraffin-embedded human lung cancer (A), cholangiocarcinorna (B), lymph node (C) and esophagus (D) tissues with DAB staining using CD73 Monoclonal Antibody.