

GPI Monoclonal Antibody

Catalog No :	YM0312
Reactivity :	Human
Applications :	WB;IHC;IF;ELISA
Target :	GPI
Fields :	>>Glycolysis / Gluconeogenesis;>>Pentose phosphate pathway;>>Starch and sucrose metabolism;>>Amino sugar and nucleotide sugar metabolism;>>Metabolic pathways;>>Carbon metabolism;>>Biosynthesis of nucleotide sugars
Gene Name :	GPI
Protein Name :	Glucose-6-phosphate isomerase
Human Gene Id :	2821
Human Swiss Prot No :	P06744
Mouse Swiss Prot No :	P06745
Immunogen :	Purified recombinant fragment of human GPI expressed in E. Coli.
Specificity :	GPI Monoclonal Antibody detects endogenous levels of GPI protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Monoclonal, Mouse
Dilution :	WB 1:500 - 1:2000. IHC 1:200 - 1:1000. IF 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other applications.
Purification :	Affinity purification
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight : 63kD

Cell Pathway : Glycolysis / Gluconeogenesis;Pentose phosphate pathway;Starch and sucrose metabolism;Amino sugar and nucleotide sugar metabolism;

P References :

1. Biochem Biophys Res Commun. 2004 Oct 15;323(2):518-22.
2. Biochem Biophys Res Commun. 2006 Oct 20;349(2):838-45.
3. Hum Mutat. 2006 Nov;27(11):1159.
4. Leuk Lymphoma. 2006 Oct;47(10):22

Background :

This gene encodes a member of the glucose phosphate isomerase protein family. The encoded protein has been identified as a moonlighting protein based on its ability to perform mechanistically distinct functions. In the cytoplasm, the gene product functions as a glycolytic enzyme (glucose-6-phosphate isomerase) that interconverts glucose-6-phosphate and fructose-6-phosphate. Extracellularly, the encoded protein (also referred to as neuroleukin) functions as a neurotrophic factor that promotes survival of skeletal motor neurons and sensory neurons, and as a lymphokine that induces immunoglobulin secretion. The encoded protein is also referred to as autocrine motility factor based on an additional function as a tumor-secreted cytokine and angiogenic factor. Defects in this gene are the cause of nonspherocytic hemolytic anemia and a severe enzyme deficiency can be associated with hydrops fetalis

Function :

catalytic activity:D-glucose 6-phosphate = D-fructose 6-phosphate.,disease:Defects in GPI are a cause of hereditary nonspherocytic hemolytic anemia (HA) [MIM:172400]. Severe GPI deficiency can be associated with hydrops fetalis, immediate neonatal death and neurological impairment.,function:Besides it's role as a glycolytic enzyme, mammalian GPI can function as a tumor-secreted cytokine and an angiogenic factor (AMF) that stimulates endothelial cell motility. GPI is also a neurotrophic factor (Neuroleukin) for spinal and sensory neurons.,online information:Phosphoglucose isomerase entry,online information:The Singapore human mutation and polymorphism database,pathway:Carbohydrate degradation; glycolysis; D-glyceraldehyde 3-phosphate and glycerone phosphate from D-glucose: step 2/4.,PTM:Phosphorylation at Ser-185 by CK2 has been shown to decrease enzymatic activity and may contribute to s

Subcellular Location : Cytoplasm . Secreted .


Expression : B-cell lymphoma,Brain,Skin,Spleen,Testis,

Sort : 6971


No4 : 1

Host : Mouse


Products Images



Western Blot analysis using GPI Monoclonal Antibody against HepG2 (1), SMMC-7721 (2) cell lysate and rat liver tissues lysate (3).



Immunohistochemistry analysis of paraffin-embedded human kidney tissues with AEC staining using GPI Monoclonal Antibody.



Confocal immunofluorescence analysis of L-02 cells using GPI Monoclonal Antibody (green). Red: Actin filaments have been labeled with DY-554 phalloidin. Blue: DRAQ5 fluorescent DNA dye.