

SNAI 1 Monoclonal Antibody

Catalog No :	YM0589
Reactivity :	Human
Applications :	WB;ELISA
Target :	SNAI1
Fields :	>>Adherens junction
Gene Name :	SNAI1
Protein Name :	Zinc finger protein SNAI1 (snail)
Human Gene Id :	6615
Human Swiss Prot No :	O95863
Mouse Swiss Prot No :	Q02085
Immunogen :	Purified recombinant fragment of human SNAI 1 expressed in E. Coli.
Specificity :	SNAI 1 Monoclonal Antibody detects endogenous levels of SNAI 1 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Monoclonal, Mouse
Dilution :	WB 1:500 - 1:2000. ELISA: 1:10000. Not yet tested in other applications.
Purification :	Affinity purification
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Molecularweight :	29kD
Cell Pathway :	Adherens_Junction;

P References :

1. Exp Cell Res. 2008 Aug 1;314(13):2448-53.
2. Mol Cell Biol. 2008 Aug;28(15):4772-81.

Background :

snail family transcriptional repressor 1(SNAI1) Homo sapiens The Drosophila embryonic protein snail is a zinc finger transcriptional repressor which downregulates the expression of ectodermal genes within the mesoderm. The nuclear protein encoded by this gene is structurally similar to the Drosophila snail protein, and is also thought to be critical for mesoderm formation in the developing embryo. At least two variants of a similar processed pseudogene have been found on chromosome 2. [provided by RefSeq, Jul 2008],

Function :

function:Seems to be involved in embryonic mesoderm formation. Binds to 3 E-boxes of the E-cadherin gene promoter and represses its transcription.,similarity:Belongs to the snail C2H2-type zinc-finger protein family.,similarity:Contains 4 C2H2-type zinc fingers.,tissue specificity:Expressed in a variety of tissues with the highest expression in kidney.,

Subcellular Location :

Nucleus . Cytoplasm . Once phosphorylated (probably on Ser-107, Ser-111, Ser-115 and Ser-119) it is exported from the nucleus to the cytoplasm where subsequent phosphorylation of the destruction motif and ubiquitination involving BTRC occurs. .

Expression :

Expressed in a variety of tissues with the highest expression in kidney. Expressed in mesenchymal and epithelial cell lines.


Sort :

16463

No4 :

1

Products Images



Western Blot analysis using SNAI 1 Monoclonal Antibody against NTERA-2 cell lysate.

