

**PRMT1 Monoclonal Antibody**


<b>Catalog No :</b>	YM1084
<b>Reactivity :</b>	Human;Mouse;Rat;Dog;Rabbit
<b>Applications :</b>	WB
<b>Target :</b>	PRMT1
<b>Fields :</b>	>>FoxO signaling pathway;>>Glucagon signaling pathway
<b>Gene Name :</b>	PRMT1
<b>Protein Name :</b>	Protein arginine N-methyltransferase 1
<b>Human Gene Id :</b>	3276
<b>Human Swiss Prot No :</b>	Q99873
<b>Mouse Gene Id :</b>	15469
<b>Mouse Swiss Prot No :</b>	Q9JIF0
<b>Rat Gene Id :</b>	60421
<b>Rat Swiss Prot No :</b>	Q63009
<b>Immunogen :</b>	Purified recombinant human PRMT1 protein fragments expressed in E.coli.
<b>Specificity :</b>	PRMT1 Monoclonal Antibody detects endogenous levels of PRMT1 protein.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Monoclonal, Mouse
<b>Dilution :</b>	WB 1:1000 - 1:2000. Not yet tested in other applications.
<b>Purification :</b>	Affinity purification

---

<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Molecularweight :</b>	42kD
<b>Background :</b>	<p>This gene encodes a member of the protein arginine N-methyltransferase (PRMT) family. Post-translational modification of target proteins by PRMTs plays an important regulatory role in many biological processes, whereby PRMTs methylate arginine residues by transferring methyl groups from S-adenosyl-L-methionine to terminal guanidino nitrogen atoms. The encoded protein is a type I PRMT and is responsible for the majority of cellular arginine methylation activity. Increased expression of this gene may play a role in many types of cancer. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene, and a pseudogene of this gene is located on the long arm of chromosome 5. [provided by RefSeq, Dec 2011],</p>
<b>Function :</b>	<p>enzyme regulation:By BTG1, BTG2 and ILF3.,function:Methylates (mono and asymmetric dimethylation) the guanidino nitrogens of arginyl residues present in a glycine and arginine-rich domain (may methylate HNRNPA1 and histones). Methylates SUPT5H and EWS.,similarity:Belongs to the protein arginine N-methyltransferase family.,subunit:Homodimer and heterodimer with PRMT8. The dimer can then associate to form a homohexamer. Interacts with ILF3, BTG1, BTG2, SUPT5H and interferon-alpha/beta receptor 1. Interacts with NFATC2IP.,</p>
<b>Subcellular Location :</b>	<p>Nucleus . Nucleus, nucleoplasm . Cytoplasm . Cytoplasm, cytosol . Mostly found in the cytoplasm. Colocalizes with CHTOP within the nucleus. Low levels detected also in the chromatin fraction (By similarity). .</p>
<b>Expression :</b>	<p>Widely expressed (PubMed:11097842). Expressed strongly in colorectal cancer cells (at protein level) (PubMed:28040436). Expressed strongly in colorectal cancer tissues compared to wild-type colon samples (at protein level) (PubMed:28040436). Expressed strongly in colorectal cancer tissues compared to wild-type colon samples (PubMed:28040436).</p>
<b>Sort :</b>	13019
<b>No4 :</b>	1
<b>Host :</b>	Mouse
<b>Modifications :</b>	Unmodified

---

## Products Images



Western Blot analysis using PRMT1 Monoclonal Antibody against HeLa, A549 cell lysate.