

Pax-4 Monoclonal Antibody

| | |
|------------------------------|--|
| Catalog No : | YM0507 |
| Reactivity : | Human |
| Applications : | WB;ELISA |
| Target : | Pax-4 |
| Fields : | >>Maturity onset diabetes of the young |
| Gene Name : | PAX4 |
| Protein Name : | Paired box protein Pax-4 |
| Human Gene Id : | 5078 |
| Human Swiss Prot No : | O43316 |
| Mouse Swiss Prot No : | P32115 |
| Immunogen : | Purified recombinant fragment of human Pax-4 expressed in E. Coli. |
| Specificity : | Pax-4 Monoclonal Antibody detects endogenous levels of Pax-4 protein. |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source : | Monoclonal, Mouse |
| Dilution : | WB 1:500 - 1:2000. ELISA: 1:10000. Not yet tested in other applications. |
| Purification : | Affinity purification |
| Storage Stability : | -15°C to -25°C/1 year(Do not lower than -25°C) |
| Molecularweight : | 38kD |
| Cell Pathway : | Maturity onset diabetes of the young; |

P References :

1. PLoS One. 2008 Mar 12;3(3):e1783.
2. Diabetes Res Clin Pract. 2008 Sep;81(3):365-9.

Background :

This gene is a member of the paired box (PAX) family of transcription factors. Members of this gene family typically contain a paired box domain, an octapeptide, and a paired-type homeodomain. These genes play critical roles during fetal development and cancer growth. The paired box 4 gene is involved in pancreatic islet development and mouse studies have demonstrated a role for this gene in differentiation of insulin-producing beta cells. [provided by RefSeq, Jul 2008],

Function :

disease:Defects in PAX4 are a cause of noninsulin-dependent diabetes mellitus (NIDDM) [MIM:125853]; also known as diabetes mellitus type 2 or maturity-onset diabetes. NIDDM is characterized by an autosomal dominant mode of inheritance, onset during adulthood and insulin resistance.,disease:Defects in PAX4 are the cause of maturity-onset diabetes of the young type 9 (MODY9) [MIM:612225]. MODY [MIM:606391] is a form of diabetes mellitus characterized by an autosomal dominant mode of inheritance, age of onset of 25 years or younger and a primary defect in insulin secretion.,disease:Genetic variations in PAX4 are associated with susceptibility to insulin-dependent diabetes mellitus (IDDM) [MIM:222100]. IDDM normally starts in childhood or adolescence and is caused by the body's own immune system which destroys the insulin-producing beta cells in the pancreas. Classical features are polydipsi

Subcellular Location :

Nucleus.

Expression :

Colon,Insulinoma,PCR rescued clones,Placenta,

Sort :

11649

No4 :

1


Host :

Mouse

Modifications :

Unmodified

Products Images



Western Blot analysis using Pax-4 Monoclonal Antibody against HEK293 (1) and PAX4-hlgGfc transfected HEK293 (2) cell lysate.

