


GRK2 mouse mAb

Catalog No :	YM1293
Reactivity :	Human
Applications :	WB
Target :	GRK 2
Fields :	>>Chemokine signaling pathway;>>Endocytosis;>>Hedgehog signaling pathway;>>Glutamatergic synapse;>>Olfactory transduction;>>Morphine addiction
Gene Name :	adrbk1
Human Gene Id :	156
Human Swiss Prot No :	P25098
Mouse Swiss Prot No :	Q99MK8
Immunogen :	Purified recombinant human GRK2 protein fragments expressed in E.coli.
Specificity :	This antibody detects endogenous levels of GRK2 and does not cross-react with related proteins.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Monoclonal, Mouse
Dilution :	wb 1:500
Purification :	The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band :	80kD
Cell Pathway :	Chemokine;Endocytosis;
Background :	<p>The product of this gene phosphorylates the beta-2-adrenergic receptor and appears to mediate agonist-specific desensitization observed at high agonist concentrations. This protein is an ubiquitous cytosolic enzyme that specifically phosphorylates the activated form of the beta-adrenergic and related G-protein-coupled receptors. Abnormal coupling of beta-adrenergic receptor to G protein is involved in the pathogenesis of the failing heart. [provided by RefSeq, Jul 2008],</p>
Function :	<p>catalytic activity:ATP + [beta-adrenergic receptor] = ADP + [beta-adrenergic receptor] phosphate.,catalytic activity:ATP + a protein = ADP + a phosphoprotein.,function:Specifically phosphorylates the agonist-occupied form of the beta-adrenergic and closely related receptors, probably inducing a desensitization of them.,online information:Beta adrenergic receptor kinase entry,similarity:Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. GPRK subfamily.,similarity:Contains 1 AGC-kinase C-terminal domain.,similarity:Contains 1 PH domain.,similarity:Contains 1 protein kinase domain.,similarity:Contains 1 RGS domain.,subunit:Interacts with GIT1 (By similarity). Interacts with, and phosphorylates chemokine-stimulated CCR5.,tissue specificity:Expressed in peripheral blood leukocytes.,</p>
Subcellular Location :	Cytoplasm . Cell membrane . Cell junction, synapse, postsynapse . Cell junction, synapse, presynapse .
Expression :	Expressed in peripheral blood leukocytes.
Sort :	7134
No4 :	1
Host :	Mouse
Modifications :	Unmodified

Products Images



Western blot detection of GRK2 in Ramos and THP-1 cell lysates using GRK2 mouse mAb (1:500 diluted), with super ECL. Predicted band size: 80KDa. Observed band size: 80KDa. Exposure time: 20S.