

OLIG2 (PTR1315) mouse mAb

Catalog No :	YM1379
Reactivity :	Human;Mouse;
Applications :	WB;IF;ELISA
Target :	OLIG2
Gene Name :	olig2
Human Gene Id :	10215
Human Swiss Prot No :	Q13516
Mouse Swiss Prot No :	Q9EQW6
Immunogen :	Synthesized peptide derived from human protein.AA range:1-100
Specificity :	This antibody detects endogenous levels of OLIG2 protein.
Formulation :	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Source :	Mouse, Monoclonal/IgG
Dilution :	WB 1:500-2000. IF 1:100-500. ELISA 1:1000-5000
Purification :	Protein G
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Molecularweight :	32kD
Observed Band :	32kD
Background :	This gene encodes a basic helix-loop-helix transcription factor which is

expressed in oligodendroglial tumors of the brain. The protein is an essential regulator of ventral neuroectodermal progenitor cell fate. The gene is involved in a chromosomal translocation t(14;21)(q11.2;q22) associated with T-cell acute lymphoblastic leukemia. Its chromosomal location is within a region of chromosome 21 which has been suggested to play a role in learning deficits associated with Down syndrome. [provided by RefSeq, Jul 2008],

Function :

disease:A chromosomal aberration involving OLIG2 may be a cause of a form of T-cell acute lymphoblastic leukemia (T-ALL). Translocation t(14;21)(q11.2;q22) with TCRA.,domain:The bHLH is essential for interaction with NKX2-2.,function:Required for oligodendrocyte and motor neuron specification in the spinal cord, as well as for the development of somatic motor neurons in the hindbrain. Cooperates with OLIG1 to establish the pMN domain of the embryonic neural tube. Antagonist of V2 interneuron and of NKX2-2-induced V3 interneuron development.,induction:By SHH. Also induced by NKX6-1 in the developing spinal cord, but not in the rostral hindbrain.,similarity:Contains 1 basic helix-loop-helix (bHLH) domain.,subcellular location:The NLS contained in the bHLH domain could be masked in the native form and translocation to the nucleus could be mediated by interaction either with class E bHLH par

Subcellular Location :

Nuclear

Expression :

Expressed in the brain, in oligodendrocytes. Strongly expressed in oligodendrogliomas, while expression is weak to moderate in astrocytomas. Expression in glioblastomas highly variable.

Sort :

11290

No4 :

1


Host :

Mouse

Modifications :

Unmodified

Products Images



Western blot detection of OLIG2 in CHO-K1 cell lysate(B) and CHO-K1 transfected by EGFP-OLIG2 fragment(A) cell lysate using OLIG2 mouse mAb (1:1000 diluted).