

**Cyclophilin B Monoclonal Antibody(2B10), FITC Conjugated**

|                              |  |
|------------------------------|--|
| <b>Catalog No :</b>          | YM2026   |
| <b>Reactivity :</b>          | Human;Rat;Mouse  |
| <b>Applications :</b>        | IF;WB;IHC;   |
| <b>Target :</b>              | Cyclophilin B  |
| <b>Gene Name :</b>           | PPIB   |
| <b>Protein Name :</b>        | Peptidyl-prolyl cis-trans isomerase B (PPIase B) (EC 5.2.1.8) (CYP-S1) (Cyclophilin B) (Rotamase B) (S-cyclophilin) (SCYLP)  |
| <b>Human Gene Id :</b>       | 5479   |
| <b>Human Swiss Prot No :</b> | P23284   |
| <b>Specificity :</b>         | Cyclophilin B Monoclonal Antibody(2B10) FITC conjugated specially designed for your WB or IHC analysis.  |
| <b>Formulation :</b>         | Liquid in PBS, pH 7.4, containing 0.02% sodium azide as preservative and 50% Glycerol.   |
| <b>Source :</b>              | Monoclonal, Mouse IgG  |
| <b>Dilution :</b>            | Optimal working dilutions should be determined experimentally by the investigator. Suggested starting dilutions are IF (1:250 - 1:2000), FCM (1:250 - 1:2000)  |
| <b>Purification :</b>        | The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.   |
| <b>Concentration :</b>       | 1mg/ml   |
| <b>Storage Stability :</b>   | Stable for one year at -15°C to -25°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid repeated freezi |

The protein encoded by this gene is a cyclosporine-binding protein and is mainly

|                               |   |
|-------------------------------|---|
| <b>Background :</b>           | located within the endoplasmic reticulum. It is associated with the secretory pathway and released in biological fluids. This protein can bind to cells derived from T- and B-lymphocytes, and may regulate cyclosporine A-mediated immunosuppression. Variants have been identified in this protein that give rise to recessive forms of osteogenesis imperfecta. [provided by RefSeq, Oct 2009],  |
| <b>Function :</b>             | catalytic activity:Peptidylproline (omega=180) = peptidylproline (omega=0).,caution:It is uncertain whether Met-1 or Met-9 is the initiator.,enzyme regulation:Cyclosporin A (CsA) inhibits CYPB.,function:PPlases accelerate the folding of proteins. It catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides.,similarity:Belongs to the cyclophilin-type PPlase family. PPlase B subfamily.,similarity:Contains 1 PPlase cyclophilin-type domain.,subcellular location:Identified by mass spectrometry in melanosome fractions from stage I to stage IV., |
| <b>Subcellular Location :</b> | Virion . (Microbial infection).; Endoplasmic reticulum lumen . Melanosome . Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:17081065). .   |
| <b>Expression :</b>           | Brain,Fetal brain cortex,Prostate,Skin,   |
| <b>Sort :</b>                 | 4757  |
| <b>No4 :</b>                  | 1   |
| <b>Host :</b>                 | Mouse   |
| <b>Modifications :</b>        | Unmodified  |
| <b>Conjugate :</b>            | FITC  |

---

## Products Images