

F۱	/n	m	οι	ıse	m	Ab)

YM1307 Catalog No:

Human; Monkey; Rat; Mouse Reactivity:

Applications: WB;ICC

Target: Fyn

Fields: >>Sphingolipid signaling pathway;>>Phospholipase D signaling

pathway:>>Axon guidance:>>Osteoclast differentiation:>>Focal

adhesion;>>Adherens junction;>>Platelet activation;>>Natural killer cell mediated

cytotoxicity;>>T cell receptor signaling pathway;>>Fc epsilon RI signaling pathway;>>Cholinergic synapse;>>Prion disease;>>Pathogenic Escherichia coli

infection;>>Viral myocarditis

Gene Name: fyn

Human Gene Id: 2534

Human Swiss Prot

No:

Mouse Swiss Prot

No:

Purified recombinant human Fyn protein fragments expressed in E.coli. Immunogen:

Specificity: This antibody detects endogenous levels of Fyn and does not cross-react with

related proteins.

P06241

P39688

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Monoclonal, Mouse Source:

Dilution: wb 1:500 icc 1:50

Purification: The antibody was affinity-purified from mouse ascites by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

1/3



Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 59kD

Cell Pathway: Axon guidance; Focal adhesion; Adherens_Junction; Natural killer cell mediated

cytotoxicity; T Cell Receptor; Fc epsilon RI; Prion diseases; Pathogenic

Escherichia coli infection; Viral myocarditis;

Background: This gene is a member of the protein-tyrosine kinase oncogene family. It

encodes a membrane-associated tyrosine kinase that has been implicated in the

control of cell growth. The protein associates with the p85 subunit of phosphatidylinositol 3-kinase and interacts with the fyn-binding protein.

Alternatively spliced transcript variants encoding distinct isoforms exist. [provided

by RefSeq, Jul 2008],

Function : catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine

phosphate.,cofactor:Manganese.,enzyme regulation:Inhibited by phosphorylation of Tyr-531 by leukocyte common antigen and activated by dephosphorylation of

this site., function: Implicated in the control of cell growth. Plays a role in the regulation of intracellular calcium levels, with isoform 2 showing the greater ability

to mobilize cytoplasmic calcium in comparison to isoform 1. Required in brain development and mature brain function with important roles in the regulation of

axon growth, axon guidance, and neurite extension. Blocks axon outgrowth and attraction induced by NTN1 by phosphorylating its receptor

DDC., similarity:Belongs to the protein kinase superfamily. Tyr protein kinase

family., similarity: Belongs to the protein kinase superfamily. Tyr protein kinase

family. SRC subfamily., similarity: Contains 1

Subcellular Cytoplasm. Nucleus. Cell membrane. Present and active in lipid rafts.

Location: Palmitoylation is crucial for proper trafficking.

Expression: Isoform 1 is highly expressed in the brain. Isoform 2 is expressed in cells of

hemopoietic lineages, especially T-lymphocytes.


Sort: 6336

No4: 1


Host: Mouse

Modifications: Unmodified

Products Images



Western blot detection of Fyn in Hela,3T3,C6,COS7,CEM,Ramos and Jurkat cell lysates using Fyn mouse mAb (1:500 diluted).Predicted band size:59KDa.Observed band size:59KDa.



Immunocytochemistry staining of HeLa cells fixed with -20°C Methanol and using Fyn mouse mAb (dilution 1:50).