

Jak1 mouse mAb

Catalog No: YM1316

Reactivity: Human;Rat

Applications: WB;IF;IP

Target: JAK1

Fields: >>EGFR tyrosine kinase inhibitor resistance;>>PI3K-Akt signaling

pathway;>>Necroptosis;>>Osteoclast differentiation;>>Signaling pathways

regulating pluripotency of stem cells;>>NOD-like receptor signaling pathway;>>JAK-STAT signaling pathway;>>Th1 and Th2 cell

differentiation;>>Th17 cell

differentiation;>>Leishmaniasis;>>Toxoplasmosis;>>Tuberculosis;>>Hepatitis C;>>Hepatitis B;>>Measles;>>Human cytomegalovirus infection;>>Influenza

A;>>Human papillomavirus infection;>>Human T-cell leukemia virus 1

infection;>>Kaposi sarcoma-associated herpesvirus infection;>>Herpes simplex

virus 1 infection;>>Epstein-Barr virus infection;>>Coronavirus disease - COVID-19;>>Pathways in cancer;>>Viral carcinogenesis;>>Pancreatic cancer;>>PD-L1 expression and PD-1 checkpoint pathway in cancer

Gene Name: jak1

Human Gene Id: 3716

Human Swiss Prot

No:

Mouse Swiss Prot

No:

Immunogen : Purified recombinant human Jak1 protein fragments expressed in E.coli.

Specificity: This antibody detects endogenous levels of Jak1 and does not cross-react with

related proteins.

P23458

P52332

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Monoclonal, Mouse

Dilution: wb 1:200-1000 icc 1:200. IF 1:50-200

1/3



Purification: The antibody was affinity-purified from mouse ascites by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 130kD

Cell Pathway : Jak_STAT;Pathways in cancer;Pancreatic cancer;

Background: This gene encodes a membrane protein that is a member of a class of protein-

tyrosine kinases (PTK) characterized by the presence of a second

phosphotransferase-related domain immediately N-terminal to the PTK domain. The encoded kinase phosphorylates STAT proteins (signal transducers and activators of transcription) and plays a key role in interferon-alpha/beta and interferon-gamma signal transduction. Alternative splicing results in multiple

transcript variants. [provided by RefSeq, Mar 2016],

Function : catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine

phosphate.,domain:Possesses two phosphotransferase domains. The second one probably contains the catalytic domain (By similarity), while the presence of slight differences suggest a different role for domain 1.,domain:The FERM domain mediates interaction with JAKMIP1.,function:Tyrosine kinase of the non-receptor type, involved in the IFN-alpha/beta/gamma signal pathway. Kinase partner for the interleukin (IL)-2 receptor.,sequence caution:Translation N-terminally extended.,similarity:Belongs to the protein kinase superfamily. Tyr

protein kinase family. JAK subfamily., similarity: Contains 1 FERM

domain., similarity: Contains 1 protein kinase domain., similarity: Contains 1 SH2

domain., subcellular location: Wholly intracellular, possibly membrane

associated., subunit: Interacts with IL31RA, JAKMIP1 and SHB., tissue specif

Subcellular Location:

Endomembrane system; Peripheral membrane protein. Wholly intracellular,

possibly membrane associated.

Expression: Expressed at higher levels in primary colon tumors than in normal colon tissue.

The expression level in metastatic colon tumors is comparable to the expression

level in normal colon tissue.

Tag: ip


Sort: 8755

No4:


Host: Mouse

Modifications: Unmodified


Products Images



Western blot analysis of extracts from C6,Ramos,Jurkat and Hela cell lysates using Jak1 mouse mAb (1:1000 diluted).Predicted band size:130KDa.Observed band size:130KDa.



Immunocytochemistry staining of HeLa cells fixed with 4% Paraformaldehyde and using anti-Jak1 mouse mAb (dilution 1:200).



Immunoprecipitation analysis of Hela cell lysates using Jak1 mouse mAb.

Hela
WB:200622-8B8 Anti-Jak1 Mouse mAb