

## Rabbit IgG mouse mAb YM1424 **Catalog No:** Rabbit **Reactivity: Applications:** WB;ELISA Target: Rabbit IgG Gene Name: rabbit igg **Human Gene Id: Human Swiss Prot** No: Rabbit IgG Immunogen: This antibody detects rabbit IgG. **Specificity:** Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Monoclonal, Mouse Source: **Dilution:** wb dilution 1:50000 **Purification:** The antibody was affinity-purified from mouse ascites by affinitychromatography using epitope-specific immunogen. **Concentration:** 1 mg/ml **Storage Stability:** -15°C to -25°C/1 year(Do not lower than -25°C) **Observed Band:** recombinant Tag:

13725

Sort:




No4:


**Host:** Mouse

Modifications: Unmodified


## **Products Images**



Western blot detection of MEK-2 in THP-1, Jurkat, Raji, 3T3 and Hela cell lysates using 310001 (MEK-2 rabbit pAb, 1:1000 diluted), then incubated with 201069 (Rabbit IgG mAb, 1:50000 diluted). Anti-mouse IgG, HRP-linked Antibody was used for detecting. Predicted band size: 44KDa. Observed band size: 44KDa.



Western blot analysis of decreasing concentrations of total rabbit IgG reduced and denatured in 2X SDS loading buffer with DTT using Mouse Anti-Rabbit IgG (Heavy-Chain Specific) mAb (1:1000 diluted).



Immunoprecipitation analysis of CHO-K1 cell lysates transfected with HA-tagged NGAL using HA-Tag Rabbit pAb (390001,1:100 diluted) (lane 2). Lane 1 contains rabbit IgG IP control lysate. All lanes are probed with HA-Tag Rabbit pAb (390001,1:1000 diluted) as the primary antibody. Panel A incubated with secondary antibodies Anti-Rabbit whole IgG, HRP-linked Antibody for detecting.Panel B incubated with Anti-Rabbit IgG (Heavy-Chain Specific) mAb(1:50000 diluted) and the Antimouse whole IgG, HRP-linked Antibody was used for detecting.