



## Negative Control for mouse IgG2b, kappa Primary Antibody

<b>Catalog No :</b>	VN0003
<b>Reactivity :</b>	Human;Mouse;Rat;Bovin;Pig;Chick;
<b>Applications :</b>	IHC;IP;IF;ELISA
<b>Human Swiss Prot No :</b>	/
<b>Formulation :</b>	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
<b>Source :</b>	Mouse, Monoclonal/IgG2b, kappa
<b>Dilution :</b>	IHC 1:200-1000. IF 1:100-500. ELISA 1:1000-5000
<b>Purification :</b>	Protein G
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Subcellular Location :</b>	No staining
<b>Sort :</b>	1
<b>No4 :</b>	1
<b>Host :</b>	Mouse
<b>Modifications :</b>	Unmodified


## Products Images




Immunohistochemical analysis of paraffin-embedded human tonsil tissue using anti-CD20 mouse mAb (ABT408, left) and mouse IgG2b, kappa monoclonal control (vn0003, right panel) .



Immunohistochemical analysis of paraffin-embedded human spleen tissue using anti-CD45 mouse mAb (left) and mouse IgG2b, kappa monoclonal control (VN003, right panel).



mmunohistochemical analysis of paraffin-embedded human prostate tissue using anti-Cytokeratin 14 mouse mAb (YM6031, left) and mouse IgG2b, kappa monoclonal control (VN003, right).



GAPDH was immunoprecipitated from Hela whole cell lysate with anti-GAPDH (PTR2304) antibody. Western blot was performed on the immunoprecipitate using anti-GAPDH antibody, and followed by the HRP-conjugated Goat anti-Mouse IgG (H + L) antibody. Lane 1 (Input) : Hela whole cell lysate Lane 2 (-) : Mouse monoclonal IgG1 (vn0003, Isotype Control) instead of GAPDH antibody in Hela whole cell lysate. Lane 3 (+) : GAPDH antibody IP in Hela whole cell lysate