


Negative Control for mouse IgG Primary Antibody


Catalog No :	VN0005
Reactivity :	Human;Mouse;Rat;Bovin;Pig;Chick;
Applications :	IHC;IP;IF;ELISA
Human Swiss Prot	/
Formulation :	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Source :	Mouse, Monoclonal/IgG
Dilution :	IHC 1:200-1000. IF 1:100-500. ELISA 1:1000-5000
Purification :	Protein G
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Subcellular	No staining
Tag :	ip
Sort :	1
No4 :	1
Host :	Mouse
Modifications :	Unmodified

Products Images





GAPDH was immunoprecipitated from SW480 whole cell lysate with anti-GAPDH (PTR2304) antibody. Western blot was performed on the immunoprecipitate using anti-GAPDH antibody, and followed by the HRP-conjugated Goat anti-Mouse IgG (H + L) antibody. Lane 1 (Input) : SW480 whole cell lysate Lane 2 (-) : Mouse monoclonal IgG (Negative Control) instead of GAPDH antibody in SW480 whole cell lysate. Lane 3 (+) : GAPDH antibody IP in SW480 whole cell lysate



Immunohistochemical analysis of paraffin-embedded human brain tissue using anti-Iba1 mouse mAb (YM4765, left) and mouse IgG, kappa monoclonal control (VN005, right).