


DDDDK-Tag(binds to flag sequence) rabbit pAb

Catalog No :	YG0004
Reactivity :	Species independent
Applications :	WB;ELISA;IP;IF
Target :	FLAG-tag
Gene Name :	Flag tag; Flag-tag, DDDDK TAG, DDDDK-TAG, DYKDDDDK tag, DYKDDDDK-tag
Immunogen :	DDDDK synthetic peptide conjugated to KLH.
Specificity :	FLAG-tag Polyclonal Antibody detects FLAG-tagged recombinant proteins or FLAG-tagged proteins overexpressed in cells.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit, IgG
Dilution :	WB 1:1000 - 1:3000. ELISA: 1:5000-20000.. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year (Do not lower than -25°C)
Background :	<p>The DYKDDDDK (FLAG) peptide has been used extensively as a general tag in expression vectors. This peptide can be expressed and detected with the protein of interest as an amino-terminal or carboxy-terminal fusion. N-terminal FLAG vectors provide an Ek cleavage site for removal of the fusion tag. The FLAG peptide is likely to be located on the surface of a fusion protein because of its hydrophilic nature. As a result, the FLAG peptide is more likely to be accessible to antibodies. A FLAG-tag can be used in many different assays that require recognition by an antibody, such as western blotting, immunocytochemistry, immunoprecipitation, flow cytometry, protein purification, and in the study of protein-protein interactions, cell ultrastructure, and protein localization and so on.</p>

Tag :	<u>orthogonal,overexpres,ip</u>
Sort :	<u>603</u>
No4 :	<u>1</u>
Host :	<u>Rabbit</u>
Modifications :	<u>Unmodified</u>

Products Images



Western Blot analysis using FLAG-tag Polyclonal Antibody against HEK293 cells transfected with vector overexpressing FLAG tag (1) and untransfected (2). Secondary antibody(catalog#:RS0002) was diluted at 1:20000