


## Goat Anti Mouse IgG(H+L)

|                            |  |
|----------------------------|--|
| <b>Catalog No :</b>        | RS0007   |
| <b>Reactivity :</b>        | Mouse  |
| <b>Applications :</b>      | WB;IHC;IF;ELISA  |
| <b>Target :</b>            | Mouse IgG(H+L)   |
| <b>Formulation :</b>       | Liquid in PBS containing, 0.5% BSA and 0.02% sodium azide.   |
| <b>Source :</b>            | Goat   |
| <b>Dilution :</b>          | WB 1:1000-10000, ELISA 1:10000-20000, IHC 1:50-100 IF 1:50-1000  |
| <b>Purification :</b>      | The antibody was affinity-purified from goat antiserum by affinity-chromatography using epitope-specific immunogen.  |
| <b>Concentration :</b>     | 4 mg/ml  |
| <b>Storage Stability :</b> | -15°C to -25°C/1 year(Do not lower than -25°C)   |
| <b>Background :</b>        | Immunoway secondary antibodies are available conjugated to enzyme, biotin or fluorophore for use in a variety of antibody-based applications including Western Blot, ImmunoHistoChemistry, ImmunoFluorescence, Flow Cytometry and ELISA. We offer high quality secondary antibodies from goat, rabbit and donkey sources for your each application. Serum adsorbed secondary antibodies are also available and are recommended for use with immunoglobulin-rich samples. |
| <b>Sort :</b>              | 6829   |
| <b>No4 :</b>               | 1  |
| <b>Host :</b>              | Goat   |
| <b>Conjugate :</b>         | unConjugated   |

## Products Images



**I** Polysaccharides from *Tetragium Hemsleyanum* Diels et Gilg ameliorated inflammatory bowel disease by rebuilding the intestinal mucosal barrier and inhibiting inflammation through the SCFA-GPR41/43 signaling pathway. INTERNATIONAL JOURNAL OF BIOLOGICAL MACROMOLECULES Fangmei Zhou