

## Lamin B1 Monoclonal Antibody(7C11), HRP Conjugated

YM2111 Catalog No:

Human;Rat;Mouse Reactivity:

**Applications:** WB;IHC;IF;IP

**Target:** Lamin B1

Fields: >>Apoptosis

Gene Name: LMNB1

**Protein Name:** Lamin-B1

**Human Gene Id:** 4001

**Human Swiss Prot** 

No:

P20700

**Specificity:** Lamin B1 Monoclonal Antibody(7C11) HRP Conjugated, specially designed for

your Western blot analysis.

Formulation: Liquid in PBS, pH 7.4, containing 0.02% sodium azide as preservative and 50%

Glycerol.

Monoclonal, Mouse IgG Source:

**Dilution:** Optimal working dilutions should be determined experimentally by the

investigator. Suggested starting dilutions are as follows: WB 1:2000-5000, IHC

1:50-300.

**Purification:** The antibody was affinity-purified from mouse ascites by affinity-

chromatography using specific immunogen.

**Concentration:** 1mg/ml

Stable for one year at -15°C to -25°C from date of shipment. For maximum **Storage Stability:** 

recovery of product, centrifuge the original vial after thawing and prior to removing

the cap. Aliquot to avoid repeated freezi

66kD **Molecularweight:** 

**Background:** lamin B1(LMNB1) Homo sapiens This gene encodes one of the two B-type lamin

> proteins and is a component of the nuclear lamina. A duplication of this gene is associated with autosomal dominant adult-onset leukodystrophy (ADLD). Alternative splicing results in multiple transcript variants. [provided by RefSeq,

Dec 2015],

**Function:** disease:Defects in LMNB1 are the cause of leukodystrophy demyelinating

> autosomal dominant adult-onset (ADLD) [MIM:169500]. ADLD is a slowly progressive and fatal demyelinating leukodystrophy, presenting in the fourth or fifth decade of life. Clinically characterized by early autonomic abnormalities, pyramidal and cerebellar dysfunction, and symmetric demyelination of the CNS. It

differs from multiple sclerosis and other demyelinating disorders in that

neuropathology shows preservation of oligodendroglia in the presence of subtotal demyelination and lack of astrogliosis., function: Lamins are components of the nuclear lamina, a fibrous layer on the nucleoplasmic side of the inner nuclear membrane, which is thought to provide a framework for the nuclear envelope and may also interact with chromatin., miscellaneous: The structural integrity of the

lamina is strictly controlled by the cell cycle

Subcellular Location:

Nucleus lamina.

Brain, Cajal-Retzius cell, Epithelium, Eye, Fetal brain cortex, Ovarian **Expression:** 

carcinoma, Placenta, Uterus,

Tag: iр

Sort: 9100

No4:

Host: Mouse

**Modifications:** Unmodified

HRP Conjugate:

## **Products Images**