

Rabbit Anti Horse IgG (H&L)-AbFluor 555

Catalog No: RS3420

Reactivity: Horse

Applications : Elisa;IF;FCM

Target: Horse IgG (H&L)

Formulation: 1 mg/ml, liquid in 0.01M Phosphate Buffered Saline, pH 7.2, containing 1%

BSA, 50% glycerol, 0.02% Sodium Azide

Source: Polyclonal, Rabbit, IgG

Dilution: IF (1:200 - 1:1000), FCM (1:100 - 1:1000), Elisa (Use at an assay dependent

concentration)

Purification: The antibody was isolated from antisera by immunoaffinity chromatography

using antigens coupled to agarose beads.

Concentration: 1mg/mL

Storage Stability: Stable for one year at -15°C to -25°C from date of shipment. For maximum

recovery of product, centrifuge the original vial after thawing and prior to removing

the cap. Aliquot to avoid repeated freezi

Background: Immunoway secondary antibodies are available conjugated to enzyme, biotin or

fluorophore for use in a variety of antibody-based applications including Western Blot, ImmunoHistoChemistry, ImmunoFluorescence, Flow Cytometry and ELISA. We offer high quality secondary antibodies from goat, rabbit and donkey sources

for your each application. Serum adsorbed secondary antibodies are also available and are recommended for use with immunoglobulin-rich samples.

Sort: 13391

No4: 1

Host: Rabbit

Conjugate: AbFluor 555



Products Images

Alexa Fluor 35	0 346/442	Blue
Alexa Fluor 40	5 401/421	Blue
Alexa Fluor 48	8 496/519	Green
Alexa Fluor 53	2 532/553	Yellow
Alexa Fluor 55	5 555/565	Yellow
Alexa Fluor 56	8 578/603	Red/Orange
Alexa Fluor 59	4 590/617	Red/Orange
Alexa Fluor 63	3 632/647	Red
Alexa Fluor 64	7 650/665	Red
Alexa Fluor 66	0 663/690	Near IR
Alexa Fluor 68	679/702	Near IR
Alexa Fluor 75	0 749/775	Near IR
Alexa Fluor 79	0 784/814	Near IR

To use the Alexa Fluors with fluorescent imagers, use a spectral line of the blue laser diode for Alexa Fluors 405, a cyan (488 nm) laser for Alexa Fluors 488, a yellow (526 nm) laser for Alexa Fluor 550 or 594, and a red (633 nm) laser for Alexa Fluor 649. The Alexa Fluor 680 and 790 fluors are compatible with laser- and filter-based infrared imaging instruments that emit in the 700 nm, and 800 nm