


PA21B Polyclonal Antibody

| | |
|------------------------------|--|
| Catalog No : | YN1055 |
| Reactivity : | Human;Rat;Mouse; |
| Applications : | WB;ELISA |
| Target : | PA21B |
| Fields : | >>Glycerophospholipid metabolism;>>Ether lipid metabolism;>>Arachidonic acid metabolism;>>Linoleic acid metabolism;>>alpha-Linolenic acid metabolism;>>Metabolic pathways;>>Ras signaling pathway;>>Vascular smooth muscle contraction;>>Pancreatic secretion;>>Fat digestion and absorption |
| Gene Name : | PLA2G1B PLA2 PLA2A PPLA2 |
| Protein Name : | Phospholipase A2 (EC 3.1.1.4) (Group IB phospholipase A2) (Phosphatidylcholine 2-acylhydrolase 1B) |
| Human Gene Id : | 5319 |
| Human Swiss Prot No : | P04054 |
| Mouse Swiss Prot No : | Q9Z0Y2 |
| Rat Swiss Prot No : | P04055 |
| Immunogen : | Synthesized peptide derived from human protein . at AA range: 70-150 |
| Specificity : | PA21B Polyclonal Antibody detects endogenous levels of protein. |
| Formulation : | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Source : | Polyclonal, Rabbit,IgG |
| Dilution : | WB 1:500-2000 ELISA 1:5000-20000 |
| Purification : | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |

| | |
|-------------------------------|---|
| Concentration : | 1 mg/ml |
| Storage Stability : | -15°C to -25°C/1 year(Do not lower than -25°C) |
| Observed Band : | 16kD |
| Cell Pathway : | Glycerophospholipid metabolism;Ether lipid metabolism;Arachidonic acid metabolism;Linoleic acid metabolism;alpha-Linolenic acid metabolism;MAPK_ERK_Growth;MAPK_G_Protein;Vascular smooth muscle contrac |
| Background : | This gene encodes a secreted member of the phospholipase A2 (PLA2) class of enzymes, which is produced by the pancreatic acinar cells. The encoded calcium-dependent enzyme catalyzes the hydrolysis of the sn-2 position of membrane glycerophospholipids to release arachidonic acid (AA) and lysophospholipids. AA is subsequently converted by downstream metabolic enzymes to several bioactive lipophilic compounds (eicosanoids), including prostaglandins (PGs) and leukotrienes (LTs). The enzyme may be involved in several physiological processes including cell contraction, cell proliferation and pathological response. [provided by RefSeq, Aug 2013], |
| Function : | catalytic activity:Phosphatidylcholine + H(2)O = 1-acylglycerophosphocholine + a carboxylate.,cofactor:Binds 1 calcium ion per subunit.,function:PA2 catalyzes the calcium-dependent hydrolysis of the 2-acyl groups in 3-sn-phosphoglycerides.,similarity:Belongs to the phospholipase A2 family., |
| Subcellular Location : | Secreted . Secreted from pancreatic acinar cells in its inactive form. |
| Expression : | Selectively expressed in pancreas, lung, liver and kidney. Also detected at lower levels in ovary and testis. |
| Sort : | 20769 |
| No4 : | 1 |
| Host : | Rabbit |
| Modifications : | Unmodified |

Products Images



Western blot analysis of lysates from A431 cells, primary antibody was diluted at 1:1000, 4° over night