

IL-33 rabbit-FC recombinant protein

Catalog No :	YD3109
Reactivity :	Human;
Purity :	>90% as determined by SDS-PAGE
Gene Name :	IL-33
Protein Name :	Interleukin-33
Sequence :	Amino acid:110-270,with rabbit FC tag.
Human Gene Id :	90865
Human Swiss Prot No :	O95760
Formulation :	Phosphate-buffered solution
Source :	Mammalian cells
Storage Stability :	-15°C to -25°C/1 year(Avoid freeze / thaw cycles)
Background :	<p>The protein encoded by this gene is a cytokine that binds to the IL1RL1/ST2 receptor. The encoded protein is involved in the maturation of Th2 cells and the activation of mast cells, basophils, eosinophils and natural killer cells. Several transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2015],</p>
Function :	<p>caution:Was originally (PubMed:12819012) thought to one of the key factors that controls the specialized post-capillary high endothelial venules (HEV) phenotype found in organized secondary lymphoid tissue and to be nuclear localized.,function:Cytokine that binds to and signals through IL1RL1/ST2 and its stimulation recruits MYD88, IRAK1, IRAK4, and TRAF6, followed by phosphorylation of MAPK3/ERK1 and/or MAPK1/ERK2, MAPK14, and MAPK8. Induces T helper type 2-associated cytokines.,PTM:Proteolytically converted to a mature form by CASP1.,similarity:Belongs to the IL-1 family. Highly divergent.,tissue specificity:Expressed at high level in high endothelial venules found in tonsils, Peyer patches and mesenteric lymph nodes. Almost undetectable in placenta.,</p>

**Subcellular
Location :**

Nucleus . Chromosome . Cytoplasm . Cytoplasmic vesicle, secretory vesicle .
Secreted . Associates with heterochromatin and mitotic chromosomes
(PubMed:17185418). The secretion is dependent on protein unfolding and
facilitated by the cargo receptor TMED10; it results in protein translocation from
the cytoplasm into the ERGIC (endoplasmic reticulum-Golgi intermediate
compartment) followed by vesicle entry and secretion (PubMed:32272059). .

Expression :

Expressed at high level in high endothelial venules found in tonsils, Peyer
patches and mesenteric lymph nodes. Almost undetectable in placenta.

Products Images