

CD55-FC recombinant protein

Catalog No: YD3086

Reactivity: Human;

Purity: >90% as determined by SDS-PAGE

Gene Name: CD55

Protein Name: Complement decay-accelerating factor (CD antigen CD55)

Sequence: Amino acid:35-353, with FC tag.

P08174

Human Gene Id: 1604

Human Swiss Prot

No:

Formulation: Phosphate-buffered solution

Source: Mammalian cells

Storage Stability: -15°C to -25°C/1 year(Avoid freeze / thaw cycles)

Function: This protein recognizes C4b and C3b fragments that condense with cell-surface

hydroxyl or amino groups when nascent C4b and C3b are locally generated during C4 and c3 activation. Interaction of daf with cell-associated C4b and C3b polypeptides interferes with their ability to catalyze the conversion of C2 and factor B to enzymatically active C2a and Bb and thereby prevents the formation of C4b2a and C3bBb, the amplification convertases of the complement cascade (PubMed:7525274). Inhibits complement activation by destabilizing and

preventing the formation of C3 and C5 convertases, which prevents complement damage (PubMed:28657829).; (Microbial infection) Acts as a receptor for Coxsackievirus A21, coxsackieviruses B1, B3 and B5.; (Microbial infection) Acts as a receptor for Human enterovirus 70 and D68 (Probable).; (Microbial infection)

Acts as a receptor for Human echoviruses 6, 7, 11, 12

Subcellular Location:

[Isoform 1]: Cell membrane; Single-pass type I membrane protein.; [Isoform 2]: Cell membrane; Lipid-anchor, GPI-anchor.; [Isoform 3]: Secreted .; [Isoform 5]: Secreted .; [Isoform 6]: Cell membrane; Lipid-anchor, GPI-

anchor .; [Isoform 7]: Cell membrane; Lipid-anchor, GPI-anchor.

1/2



Expression:

Expressed on the plasma membranes of all cell types that are in intimate contact with plasma complement proteins. It is also found on the surfaces of epithelial cells lining extracellular compartments, and variants of the molecule are present in body fluids and in extracellular matrix.

Products Images