

CD22-FC recombinant protein

| | |
|-------------------------------|--|
| Catalog No : | YD3076 |
| Reactivity : | Human; |
| Purity : | >90% as determined by SDS-PAGE |
| Gene Name : | CD22 |
| Protein Name : | B-cell receptor CD22 (B-lymphocyte cell adhesion molecule) (BL-CAM) (Sialic acid-binding Ig-like lectin 2) (Siglec-2) (T-cell surface antigen Leu-14) (CD antigen CD22) |
| Sequence : | Amino acid:20-687,with FC tag. |
| Human Gene Id : | 933 |
| Human Swiss Prot No : | P20273 |
| Formulation : | Phosphate-buffered solution |
| Source : | Mammalian cells |
| Storage Stability : | -15°C to -25°C/1 year(Avoid freeze / thaw cycles) |
| Function : | Mediates B-cell B-cell interactions. May be involved in the localization of B-cells in lymphoid tissues. Binds sialylated glycoproteins; one of which is CD45. Preferentially binds to alpha-2,6-linked sialic acid. The sialic acid recognition site can be masked by cis interactions with sialic acids on the same cell surface. Upon ligand induced tyrosine phosphorylation in the immune response seems to be involved in regulation of B-cell antigen receptor signaling. Plays a role in positive regulation through interaction with Src family tyrosine kinases and may also act as an inhibitory receptor by recruiting cytoplasmic phosphatases via their SH2 domains that block signal transduction through dephosphorylation of signaling molecules. |
| Subcellular Location : | Cell membrane; Single-pass type I membrane protein. |
| Expression : | B-lymphocytes. |

Products Images