


EAAT1 (PT0196R) PT® Rabbit mAb


Catalog No :	YM8123
Reactivity :	Human; Mouse; Rat;
Applications :	WB;IHC;IF;IP;ELISA
Target :	EAAT1
Fields :	>>Synaptic vesicle cycle;>>Glutamatergic synapse;>>Huntington disease
Gene Name :	SLC1A3
Protein Name :	Excitatory amino acid transporter 1
Human Gene Id :	6507
Human Swiss Prot No :	P43003
Mouse Swiss Prot No :	P56564
Specificity :	endogenous
Formulation :	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Source :	Monoclonal, rabbit, IgG, Kappa
Dilution :	IHC 1:200-1000,WB 1:500-2000,IF 1:200-1000,ELISA 1:5000-20000,IP 1:50-200
Purification :	Protein A
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Molecularweight :	60kD
Observed Band :	59kD

Background :	This gene encodes a member of a member of a high affinity glutamate transporter family. This gene functions in the termination of excitatory neurotransmission in central nervous system. Mutations are associated with episodic ataxia, Type 6. Alternative splicing results in multiple transcript variants.[provided by RefSeq, Feb 2014],
Function :	disease:Defects in SLC1A3 are the cause of episodic ataxia type 6 (EA6) [MIM:612656]. EA6 is characterized by episodic ataxia, seizures, migraine and alternating hemiplegia.,function:Transports L-glutamate and also L- and D-aspartate. Essential for terminating the postsynaptic action of glutamate by rapidly removing released glutamate from the synaptic cleft. Acts as a symport by cotransporting sodium.,PTM:Glycosylated.,similarity:Belongs to the sodium:dicarboxylate (SDF) symporter (TC 2.A.23) family.,tissue specificity:Highly expressed in cerebellum, but also found in frontal cortex, hippocampus and basal ganglia.,
Subcellular Location :	Cell membrane
Expression :	Detected in brain (PubMed:8218410, PubMed:7521911, PubMed:8123008). Detected at very much lower levels in heart, lung, placenta and skeletal muscle (PubMed:7521911, PubMed:8123008). Highly expressed in cerebellum, but also found in frontal cortex, hippocampus and basal ganglia (PubMed:7521911).
Tag :	hot,recombinant
Sort :	5364
No4 :	1
Host :	Rabbit
Modifications :	Unmodified


Products Images




Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-EAAT1 (PT0196R) antibody. The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1: U-87MG
Predicted band size: 60kDa Observed band size: 59kDa




Rat brain was stained with Anti-EAAT1 (PT0196R) rabbit antibody



Mouse brain was stained with Anti-EAAT1 (PT0196R) rabbit antibody



Human brain was stained with Anti-EAAT1 (PT0196R) rabbit antibody



Human glioma was stained with anti-EAAT1 (PT0196R) rabbit antibody