

## NKp80-FC recombinant protein

|                               |   |
|-------------------------------|---|
| <b>Catalog No :</b>           | YD3093  |
| <b>Reactivity :</b>           | Human;  |
| <b>Purity :</b>               | >90% as determined by SDS-PAGE  |
| <b>Gene Name :</b>            | KLRF1   |
| <b>Protein Name :</b>         | Killer cell lectin-like receptor subfamily F member 1 (Lectin-like receptor F1) (Activating coreceptor NKp80) (C-type lectin domain family 5 member C)  |
| <b>Sequence :</b>             | Amino acid:60-231, with FC tag.   |
| <b>Human Gene Id :</b>        | 51348   |
| <b>Human Swiss Prot No :</b>  | Q9NZS2  |
| <b>Formulation :</b>          | Phosphate-buffered solution   |
| <b>Source :</b>               | Mammalian cells   |
| <b>Storage Stability :</b>    | -15°C to -25°C/1 year(Avoid freeze / thaw cycles)   |
| <b>Function :</b>             | Involved in the natural killer (NK)-mediated cytolysis of PHA-induced lymphoblasts.   |
| <b>Subcellular Location :</b> | Membrane ; Single-pass type II membrane protein .   |
| <b>Expression :</b>           | Strongly expressed in peripheral blood leukocytes and spleen, with weaker expression in lymph node and adult liver, and no expression detected in bone marrow, thymus, and fetal liver. Not expressed in brain, heart, placenta, lung, kidney, skeletal muscle, and pancreas. Within peripheral blood leukocyte and immunocyte cell lines, expression was predominant in NK cells but was also detected in monocytes. |

## Products Images