

Caspase-9 Polyclonal Antibody

Catalog No: YT0662

Reactivity: Human; Mouse; Rat

Applications: WB;IHC;IF;ELISA

Target: Caspase-9

Fields: >>Platinum drug resistance;>>p53 signaling pathway;>>Pl3K-Akt signaling

pathway;>>Apoptosis;>>Apoptosis - multiple species;>>VEGF signaling

pathway;>>Thyroid hormone signaling pathway;>>Alzheimer

disease;>>Parkinson disease;>>Amyotrophic lateral sclerosis;>>Huntington disease;>>Prion disease;>>Pathways of neurodegeneration - multiple

diseases;>>Pathogenic Escherichia coli

infection;>>Legionellosis;>>Toxoplasmosis;>>Tuberculosis;>>Hepatitis C;>>Hepatitis B;>>Measles;>>Human cytomegalovirus infection;>>Influenza A;>>Kaposi sarcoma-associated herpesvirus infection;>>Herpes simplex virus 1 infection;>>Epstein-Barr virus infection;>>Human immunodeficiency virus 1

infection;>>Pathways in cancer;>>Colorectal cancer;>>Pancreatic

cancer;>>Endometrial cancer;>>Prostate cancer;>>Small cell lung cancer;>>Non-

small cell lung cancer;>>Viral myocarditis;>>Lipid and atherosclerosis

Gene Name: CASP9

Protein Name: Caspase9

Human Gene Id: 842

Human Swiss Prot

P55211

No:

Rat Gene Id: 58918

Rat Swiss Prot No: Q9JHK1

Immunogen: The antiserum was produced against synthesized peptide derived from human

Caspase 9. AA range:91-140

Specificity: Caspase-9 Polyclonal Antibody detects endogenous levels of Caspase-9

protein.



Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution: WB 1:500 - 1:2000. IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:5000. Not

yet tested in other applications.

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 46kD

Cell Pathway: p53;Apoptosis_Inhibition;Apoptosis_Mitochondrial;Apoptosis_Overview;VEGF;

Alzheimer's disease; Parkinson's disease; Amyotrophic lateral sclerosis

(ALS); Huntington's disease; Pathways in cancer; Colorectal

Background : This gene encodes a member of the cysteine-aspartic acid protease (caspase)

family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. This protein can

undergo autoproteolytic processing and activation by the apoptosome, a protein complex of cytochrome c and the apoptotic peptidase activating factor 1; this step is thought to be one of the earliest in the caspase activation cascade. This protein

is thought to play a central role in apoptosis and to be a tumor suppressor. Alternative splicing results in multiple transcript variants. [provided by RefSeq,

May 2013],

Function: catalytic activity: Strict requirement for an Asp residue at position P1 and with a

marked preference for His at position P2. It has a preferred cleavage sequence of Leu-Gly-His-Asp-|-Xaa.,function:Involved in the activation cascade of caspases responsible for apoptosis execution. Binding of caspase-9 to Apaf-1 leads to activation of the protease which then cleaves and activates caspase-3.

Proteolytically cleaves poly(ADP-ribose) polymerase (PARP).,function:Isoform 2

Totally lically cleaves poly (ABT Tibose) polymerase (17411)., ranciformsolorm

lacks activity is an dominant-negative inhibitor of caspase-9.,online

information:Caspase-9 entry,PTM:Cleavages at Asp-315 by granzyme B and at Asp-330 by caspase-3 generate the two active subunits. Caspase-8 and -10 can also be involved in these processing events.,similarity:Belongs to the peptidase C14A family.,similarity:Contains 1 CARD domain.,subunit:Heterotetramer that

consists of two anti-parallel arranged heterodimers

Subcellular nucleus, mitochondrion, cytosol, apoptosome,

2/3



Expatission: Ubiquitous, with highest expression in the heart, moderate expression in liver,

skeletal muscle, and pancreas. Low levels in all other tissues. Within the heart,

specifically expressed in myocytes.

Sort: 2

No3: ab32539

No4: 1

Host: Rabbit

Modifications: Unmodified

Products Images

3/3