

Adrenocorticotropin(ACTH) (ABT181R) rabbit mAb

Catalog No :	YM7007
Reactivity :	Human;
Applications :	IHC; ELISA
Target :	Adrenocorticotropin
Fields :	>>cAMP signaling pathway;>>Neuroactive ligand-receptor interaction;>>Estrogen signaling pathway;>>Melanogenesis;>>Adipocytokine signaling pathway;>>Aldosterone synthesis and secretion;>>Cortisol synthesis and secretion;>>Cushing syndrome
Gene Name :	POMC
Protein Name :	Pro-opiomelanocortin (POMC) (Corticotropin-lipotropin) [Cleaved into: NPP; Melanotropin gamma (Gamma-MSH); Potential peptide; Corticotropin (Adrenocorticotropic hormone) (ACTH); Melanotropin alpha (AI
Human Gene Id :	5443
Human Swiss Prot No :	P01189
Immunogen :	Synthesized peptide derived from human Adrenocorticotropin(ACTH) AA range:100-200
Specificity :	This antibody detects endogenous levels of Adrenocorticotropin
Formulation :	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Source :	Monoclonal, Rabbit IgG1, Kappa
Dilution :	IHC 1:100-500, ELISA 1:5000-20000
Purification :	Recombinant Expression and Affinity purified
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Cell Pathway :	Melanogenesis;Adipocytokine;
Background :	<p>This gene encodes a preproprotein that undergoes extensive, tissue-specific, post-translational processing via cleavage by subtilisin-like enzymes known as prohormone convertases. There are eight potential cleavage sites within the preproprotein and, depending on tissue type and the available convertases, processing may yield as many as ten biologically active peptides involved in diverse cellular functions. The encoded protein is synthesized mainly in corticotroph cells of the anterior pituitary where four cleavage sites are used; adrenocorticotrophin, essential for normal steroidogenesis and the maintenance of normal adrenal weight, and lipotropin beta are the major end products. In other tissues, including the hypothalamus, placenta, and epithelium, all cleavage sites may be used, giving rise to peptides with roles in pain and energy homeostasis, melanocyte stimulation, and immune modulation. The</p>
Function :	<p>disease:Defects in POMC are the cause of pro-opiomelanocortin deficiency [MIM:609734]. Affected individuals present early-onset obesity, adrenal insufficiency and red hair.,disease:Defects in POMC may be associated with susceptibility to obesity [MIM:601665].,function:ACTH stimulates the adrenal glands to release cortisol.,function:Beta-endorphin and Met-enkephalin are endogenous opiates.,function:MSH (melanocyte-stimulating hormone) increases the pigmentation of skin by increasing melanin production in melanocytes.,online information:Melanocyte-stimulating hormone entry,PTM:O-glycosylated; reducing sugar is probably N-acetylgalactosamine.,PTM:Specific enzymatic cleavages at paired basic residues yield the different active peptides.,similarity:Belongs to the POMC family.,tissue specificity:ACTH and MSH are produced by the pituitary gland.,</p>
Subcellular Location :	Cytoplasmic
Expression :	ACTH and MSH are produced by the pituitary gland.
Tag :	hot,recombinant
Sort :	800
No4 :	1
Host :	Rabbit
Modifications :	Unmodified

Products Images