

**MAGE-C1 (PT0265R) rabbit mAb**

<b>Catalog No :</b>	YM7147
<b>Reactivity :</b>	Human;
<b>Applications :</b>	IHC;WB; ELISA
<b>Target :</b>	MAGE-C1
<b>Gene Name :</b>	MAGEC1
<b>Protein Name :</b>	MAGE-C1
<b>Human Gene Id :</b>	9947
<b>Human Swiss Prot No :</b>	O60732
<b>Immunogen :</b>	Synthesized peptide derived from human MAGE-C1 AA range:700-800
<b>Specificity :</b>	This antibody detects endogenous levels of MAGE-C1
<b>Formulation :</b>	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
<b>Source :</b>	Monoclonal, Rabbit IgG1, Kappa
<b>Dilution :</b>	IHC 1:100-500, WB 1:500-1000, ELISA 1:5000-20000
<b>Purification :</b>	Recombinant Expression and Affinity purified
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Molecularweight :</b>	124kD
<b>Background :</b>	This gene is a member of the melanoma antigen gene (MAGE) family. The proteins of this family are tumor-specific antigens that can be recognized by autologous cytolytic T lymphocytes. This protein contains a large number of unique short repetitive sequences in front of the MAGE-homologous sequence, and therefore is about 800 aa longer than the other MAGE proteins. [provided by RefSeq, Jul 2008],

<b>Function :</b>	similarity:Contains 1 MAGE domain.,tissue specificity:Expressed in testis and in tumors of a wide variety of histologic types.,
<b>Subcellular Location :</b>	Cytoplasmic
<b>Expression :</b>	Expressed in testis and in tumors of a wide variety of histologic types.
<b>Tag :</b>	hot,recombinant
<b>Sort :</b>	800
<b>No4 :</b>	1
<b>Host :</b>	Rabbit
<b>Modifications :</b>	Unmodified

## Products Images