


DHSO rabbit pAb


Catalog No :	YN3626
Reactivity :	Human;Mouse;Rat
Applications :	WB;IP
Target :	DHSO
Fields :	>>Pentose and glucuronate interconversions;>>Fructose and mannose metabolism;>>Metabolic pathways
Gene Name :	SORD
Protein Name :	DHSO
Human Gene Id :	6652
Human Swiss Prot No :	Q00796
Mouse Gene Id :	20322
Mouse Swiss Prot No :	Q64442
Rat Gene Id :	24788
Rat Swiss Prot No :	P27867
Immunogen :	Synthesized peptide derived from human DHSO
Specificity :	This antibody detects endogenous levels of DHSO at Human/Mouse/Rat
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1[?]500-2000 IP 1:200

Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Molecularweight :	39kD
Background :	Sorbitol dehydrogenase (SORD; EC 1.1.1.14) catalyzes the interconversion of polyols and their corresponding ketoses, and together with aldose reductase (ALDR1; MIM 103880), makes up the sorbitol pathway that is believed to play an important role in the development of diabetic complications (summarized by Carr and Markham, 1995 [PubMed 8535074]). The first reaction of the pathway (also called the polyol pathway) is the reduction of glucose to sorbitol by ALDR1 with NADPH as the cofactor. SORD then oxidizes the sorbitol to fructose using NAD(+) cofactor.[supplied by OMIM, Jul 2010],
Function :	catalytic activity:L-iditol + NAD(+) = L-sorbose + NADH.,cofactor:Binds 1 zinc ion per subunit.,similarity:Belongs to the zinc-containing alcohol dehydrogenase family.,subunit:Homotetramer.,
Subcellular Location :	Mitochondrion membrane ; Peripheral membrane protein . Cell projection, cilium, flagellum . Associated with mitochondria of the midpiece and near the plasma membrane in the principal piece of the flagellum. Also found in the epididymosome, secreted by the epididymal epithelium and that transfers proteins from the epididymal fluid to the sperm surface. .
Expression :	Expressed in liver (PubMed:3365415). Expressed in kidney and epithelial cells of both benign and malignant prostate tissue. Expressed in epididymis (at protein level).
Tag :	ip
Sort :	18796
No4 :	1
Host :	Rabbit
Modifications :	Unmodified

Products Images



Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4° over night



1) Input: Hela Lysate 2) IP product: IP dilute 1: 200 Western blot analysis: primary antibody : 1:2000 Secondary antibody: Goat anti-Mouse IgG(RS0002), 1: 5000