

CD141 (Thrombomodulin) (ABT87R) rabbit mAb

Catalog No :	YM7034
Reactivity :	Human;Mouse;(predicted: Rat)
Applications :	IHC;WB; ELISA
Target :	CD141
Fields :	>>Complement and coagulation cascades;>>AGE-RAGE signaling pathway in diabetic complications;>>Fluid shear stress and atherosclerosis
Gene Name :	THBD
Protein Name :	CD141
Human Gene Id :	7056
Human Swiss Prot No :	P07204
Immunogen :	Synthesized peptide derived from human CD141 AA range:50-150
Specificity :	This antibody detects endogenous levels of CD141
Formulation :	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Source :	Monoclonal, Rabbit IgG1, Kappa
Dilution :	IHC 1:100-500, WB 1:500-1000, ELISA 1:5000-20000
Purification :	Recombinant Expression and Affinity purified
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Molecularweight :	60kD
Background :	The protein encoded by this intronless gene is an endothelial-specific type I membrane receptor that binds thrombin. This binding results in the activation of protein C, which degrades clotting factors Va and VIIIa and reduces the amount

of thrombin generated. Mutations in this gene are a cause of thromboembolic disease, also known as inherited thrombophilia. [provided by RefSeq, Jul 2008],

Function :

disease:Defects in THBD are the cause of thrombophilia due to thrombomodulin defect (THR-THBDD) [MIM:188040]. THR-THBDD is a hemostatic disorder characterized by a tendency to thrombosis.,function:Thrombomodulin is a specific endothelial cell receptor that forms a 1:1 stoichiometric complex with thrombin. This complex is responsible for the conversion of protein C to the activated protein C (protein Ca). Once evolved, protein Ca scissions the activated cofactors of the coagulation mechanism, factor Va and factor VIIIa, and thereby reduces the amount of thrombin generated.,online information:Thrombomodulin,online information:Thrombomodulin entry,PTM:N-glycosylated.,PTM:The iron and 2-oxoglutarate dependent 3-hydroxylation of aspartate and asparagine is (R) stereospecific within EGF domains.,similarity:Contains 1 C-type lectin domain.,similarity:Contains 6 EGF-like domains.,tissue specific

Subcellular Location :

Membranous

Expression :

Endothelial cells are unique in synthesizing thrombomodulin.

Tag :

recombinant

Sort :

800

No4 :

1

Host :

Rabbit

Modifications :

Unmodified

Products Images