

MUC4 (PT0270R) rabbit mAb

Catalog No: YM7158

Reactivity: Human;

Applications: IHC; ELISA

Target: MUC4

Gene Name: MUC4

Protein Name: MUC4

Human Gene Id: 3662

Human Swiss Prot

No:

Immunogen: Synthesized peptide derived from human MUC4 AA range:2000-2100

Specificity: This antibody detects endogenous levels of MUC4

Formulation: PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA

Source: Monoclonal, Rabbit IgG1, Kappa

Q99102

Dilution : IHC 1:100-500, ELISA 1:5000-20000

Purification: Recombinant Expression and Affinity purified

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Function: alternative products:Additional isoforms exist, developmental stage:Expressed

early in the primitive gut before respiratory and digestive epithelial cells have acquired their tissue and cell specificity. Expressed at the basal surface of the epithelium from week 14 to 26 weeks and then predominantly localized in only parietal cells. Immediately before birth, found in the cytoplasm of the mucous columnar epithelial cells. In the embryo expressed in skin, then disappears late in gestation.,function:May play a role in tumor progression. Ability to promote tumor growth may be mainly due to repression of apoptosis as opposed to proliferation. Has anti-adhesive properties. Seems to alter cellular behavior through both anti-



adhesive effects on cell-cell and cell-extracellular matrix interactions and in its ability to act as an intramembrane ligand for ERBB2. Plays an important role in cell proli

Subcellular Location:

[Mucin-4 beta chain]: Cell membrane; Single-pass membrane protein. Isoforms lacking the Cys-rich region, EGF-like domains and transmembrane region are secreted. Secretion occurs by splicing or proteolytic processing. .; [Mucin-4 alpha chain]: Cell membrane. Secreted. Forms a complex with Mucin-4 beta chain at the cell membrane. .; [Isoform 3]: Cell membrane; Single-pass membrane protein .; [Isoform 15]: Secreted .; [Isoform 17]: Cell membrane; Single-pass membrane protein.

Expression:

Expressed in the thymus, thyroid, lung, trachea, esophagus, stomach, small intestine, colon, testis, prostate, ovary, uterus, placenta, and mammary and salivary glands. Expressed in carcinomas arising from some of these epithelia, such as lung cancers, squamous cell carcinomas of the upper aerodigestive tract, mammary carcinomas, biliary tract, colon, and cervix cancers. Minimally or not expressed in the normal pancreas or chronic pancreatitis, but is highly expressed in pancreatic tumors and pancreatic tumor cell lines.

Tag: recombinant

Sort : 10335

No4: 1

Host: Rabbit

Modifications: Unmodified

Products Images

2/2